

A-level BIOLOGY (7402/3)

Paper 3

Specimen 2014

Session

Time allowed: 2 hours

Materials

For this paper you must have:

- a ruler with millimetre measurements
- a calculator.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the bottom of this page.
- Answer **all** questions in Section **A**.
- Answer one question from Section B.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 78.

Please write clearly, in block capitals, to allow character computer recognition.					
Centre number					
Surname					
Forename(s)					
Candidate signature					

	Section A Answer all questions in this section.	
1	Ecologists investigated the size of an insect population on a small island. mark-release-recapture method. To mark the insects they used a fluoresc This powder glows bright red when exposed to ultraviolet (UV) light.	
01.1	The ecologists captured insects from a number of sites on the island. Sug they decided where to take their samples.	gest how [2 marks]
01.2	Give two assumptions made when using the mark-release-recapture meth	iod. [2 marks]
	2	
01.3	Suggest the advantage of using the fluorescent powder in this experiment.	[2 marks]

	The ecologists did not release any of the insects they captured 1–5 days after release of the marked insects.
	Table 1 shows the ecologists' results.
	This source has been removed due to third-party copyright restrictions.
0 1 . 4	Calculate the number of insects on this island 1 day after release of the marked
	insects.
	Show your working.
	[2 marks]
	Answer =
01.5	The ecologists expected to obtain the same result from their calculations of the number of insects on this island on each day during the period 1–5 days after release. In fact, their estimated number increased after day 1.
	During the same period, the number of insects they caught decreased.
	The method used by the ecologists might have caused these changes.
	Use the information provided to suggest one way in which the method used by the ecologists might have caused the increase in their estimates of the size of the insect population.
	[2 marks]

02.1	Describe how you could make a temporary mount of a piece of plant tissue to observe the position of starch grains in the cells when using an optical (light) microscope. [4 marks]
	[Extra space]
	Figure 1 shows a microscopic image of a plant cell.
	This source has been removed due to third-party copyright restrictions.

02.2	Give the name and function of the structures labelled W and Z . [2 marl	ks]
	Name of W	
	Function of W	
	Name of Z	
	Function of Z	
02.3	A transmission electron microscope was used to produce the image in Figure 1 . Explain why. [2 mark]	ks]
02.4	Calculate the magnification of the image shown in Figure 1 . [1 ma	ark]
	Answer =	

3	A student investigated the effect of chewing on the digestion of starch in cooked wheat.
	He devised a laboratory model of starch digestion in the human gut. This is the method he used.
	 Volunteers chewed cooked wheat for a set time. The wheat had been cooked in boiling water. This chewed wheat was mixed with water, hydrochloric acid and a protein-digesting enzyme and left at 37 °C for 30 minutes. A buffer was then added to bring the pH to 6.0 and pancreatic amylase was added. This mixture was then left at 37 °C for 120 minutes. Samples of the mixture were removed at 0, 10, 20, 40, 60 and 120 minutes, and the concentration of reducing sugar in each sample was measured. Control experiments were carried out using cooked wheat that had been chopped up in a blender, not chewed.
03.1	What reducing sugar, or sugars, would you expect to be produced during chewing? Give a reason for your answer. [2 marks]
03.2	In this model of digestion in the human gut, what other enzyme is required for the complete digestion of starch? [1 mark]
03.3	What was the purpose of step 2, in which samples were mixed with water, hydrochloric acid and pepsin? [1 mark]

03.4	In the control experiments, cooked wheat was chopped up to copy the effect of chewing.		
	Suggest a more appropriate control experiment. Explain your suggestion. [2 marks]		
03.5	Figure 2 shows the student's results.		
	This source has been removed due to third-party copyright restrictions.		
	Explain what these results suggest about the effect of chewing on the digestion of starch in wheat. [3 marks]		

4	Metastatic melanoma (MM) is a type of skin cancer. It is caused by a faulty receptor protein in cell-surface membranes. There have been no very effective treatments for this cancer.				
	Dacarbazine is a drug that has been used to treat MM because it appears to increase survival time for some people with MM.				
	Doctors investigated the use of a new drug, called ipilimumab, to treat MM. They compared the median survival time (ST) for two groups of patients treated for MM:				
		• •	of patients who had been treated nts who had been treated with d		
		The ST is how long	a patient lives after diagnosis.		
		The doctors also red in tumours with eacl	corded the percentage of patient	ts showing a significant reduc	ction
		The total number of	patients in the investigation was	s 502.	
		Table 2 shows the c	loctors' results.		
			Table 2		
	Treati	nent	Median survival time (ST) / months	Percentage of patients showing significant reduction in tumours	
	Dacar	bazine	9.1	10.3	
	Dacarbazine and ipilimumab		11.2	15.2	
0 4 . 1 The doctors compared median survival times for patients in each group. How would you find the median survival time for a group of patients? [2 mar					narks]

In many trials of new drugs, a control group of patients is given a placebo that does not contain any drug.
The control group in this investigation had been treated with dacarbazine. Suggest why they had not been given a placebo.
[1 mark]
A journalist who read this investigation concluded that ipilimumab improved the treatment of MM.
Do the data in Table 2 support this conclusion? Give reasons for your answer. [4 marks]
[Extra space]
Question 4 continues on the next page

04.4	MM is caused by a faulty receptor protein in cell-surface membranes. Cells in MM tumours can be destroyed by the immune system.	
	Suggest why they can be destroyed by the immune system.	[3 marks]
	[Extra space]	


5	Scientists investigated the effect of a mycorrhizal fungus on the growth of pea plants with a nitrate fertiliser or an ammonium fertiliser. The fertilisers were identical, except for nitrate or ammonium. The scientists took pea seeds and sterilised their surfaces. They planted the seeds in soil that had been heated to 85 °C for 2 days before use. The soil was sand that contained no mineral ions useful to the plants.
0 5 . 1	Explain why the scientists sterilised the surfaces of the seeds and grew them in soil that had been heated to 85 °C for 2 days. [2 marks]
0 5 . 2	Explain why it was important that the soil contained no mineral ions useful to the plants. [1 mark] The pea plants were divided into four groups, A , B , C and D .
	 Group A – heat-treated mycorrhizal fungus added, nitrate fertiliser Group B – mycorrhizal fungus added, nitrate fertiliser Group C – heat-treated mycorrhizal fungus added, ammonium fertiliser Group D – mycorrhizal fungus added, ammonium fertiliser
05.3	Explain how groups A and C act as controls. [2 marks]
	Question 5 continues on the next page

0 5 . 4	After 6 weeks, the scientists removed the plants from the soil and cut the root the shoots. They dried the plant material in an oven at 90 °C for 3 days. The determined the mean dry masses of the roots and shoots of each group of personal structures what the scientists should have done during the drying process to b that all of the water had been removed from the plant samples.	ey then ea plants.
	The scientists' results are shown in Table 3 .	
	Table 3	
	This source has been removed due to third-party copyright restrictions.	

0 5 . 5	What conclusions can be drawn from the data in Table 3 about the following? [4 marks] The effects of the fungus on growth of the pea plants.
	The effects of nitrate fertiliser and ammonium fertiliser on growth of the pea plants.
	Question 5 continues on the next page

	The scientists determined the dry mass of the roots and shoots separately. The reason for this was they were interested in the ratio of shoot to root growth of pea plants. It is the shoot of the pea plant that is harvested for commercial purposes.	
0 5 . 6	investigation.	arks]
05.7	Which treatment gave the best result in commercial terms? Justify your answer. [2 m	arks]

	Section B Answer one question.	
6	Write an essay on one of the topics below.	
0 6 . 1	The importance of movement in living organisms.	[25 marks]
0 6 . 2	The importance of receptors in living organisms.	[25 marks]



-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
-	
	END OF QUESTIONS

