



# **Physics Equations Sheet**

**GCSE Physics (8463)**

**FOR USE IN JUNE 2024 ONLY**

**[Turn over]**

## HT = Higher Tier only equations

2

<b>kinetic energy = <math>0.5 \times \text{mass} \times (\text{speed})^2</math></b>	$E_k = \frac{1}{2} m v^2$
<b>elastic potential energy = <math>0.5 \times \text{spring constant} \times (\text{extension})^2</math></b>	$E_e = \frac{1}{2} k e^2$
<b>gravitational potential energy = <math>\text{mass} \times \text{gravitational field strength} \times \text{height}</math></b>	$E_p = m g h$
<b>change in thermal energy = <math>\text{mass} \times \text{specific heat capacity} \times \text{temperature change}</math></b>	$\Delta E = m c \Delta \theta$
<b>power = <math>\frac{\text{energy transferred}}{\text{time}}</math></b>	$P = \frac{E}{t}$
<b>power = <math>\frac{\text{work done}}{\text{time}}</math></b>	$P = \frac{W}{t}$
<b>efficiency = <math>\frac{\text{useful output energy transfer}}{\text{total input energy transfer}}</math></b>	

$\text{efficiency} = \frac{\text{useful power output}}{\text{total power input}}$	
$\text{charge flow} = \text{current} \times \text{time}$	$Q = I t$
$\text{potential difference} = \text{current} \times \text{resistance}$	$V = I R$
$\text{power} = \text{potential difference} \times \text{current}$	$P = V I$
$\text{power} = (\text{current})^2 \times \text{resistance}$	$P = I^2 R$
$\text{energy transferred} = \text{power} \times \text{time}$	$E = P t$
$\text{energy transferred} = \text{charge flow} \times \text{potential difference}$	$E = Q V$
$\text{density} = \frac{\text{mass}}{\text{volume}}$	$\rho = \frac{m}{V}$
$\text{thermal energy for a change of state} =$ $\text{mass} \times \text{specific latent heat}$	$E = m L$

[Turn over]

<b>For gases:</b> pressure × volume = constant	$p V = \text{constant}$
<b>weight = mass × gravitational field strength</b>	$W = m g$
<b>work done = force × distance (along the line of action of the force)</b>	$W = F s$
<b>force = spring constant × extension</b>	$F = k e$
<b>moment of a force = force × distance (normal to direction of force)</b>	$M = F d$
<b>pressure = <u>force normal to a surface</u></b> <u>area of that surface</u>	$p = \frac{F}{A}$
<b>pressure due to a column of liquid = height of column × density of liquid × gravitational field strength</b>	$p = h \rho g$
<b>distance travelled = speed × time</b>	$s = v t$
<b>acceleration = <u>change in velocity</u></b> <u>time taken</u>	$a = \frac{\Delta v}{t}$

(final velocity) <sup>2</sup> – (initial velocity) <sup>2</sup> = 2 × acceleration × distance	$v^2 - u^2 = 2 a s$
<b>resultant force</b> = mass × acceleration	$F = m a$
<b>HT momentum</b> = mass × velocity	$p = m v$
<b>HT force</b> = <u>change in momentum</u> <u>time taken</u>	$F = \frac{m \Delta v}{\Delta t}$
<b>HT period</b> = <u>1</u> <u>frequency</u>	$T = \frac{1}{f}$
<b>wave speed</b> = frequency × wavelength	$v = f \lambda$
<b>magnification</b> = <u>image height</u> <u>object height</u>	
<b>HT force on a conductor (at right angles to a magnetic field) carrying a current</b> = magnetic flux density × current × length	$F = B I l$

<b>HT</b> $\frac{\text{potential difference across primary coil}}{\text{potential difference across secondary coil}} = \frac{V_p}{V_s} = \frac{n_p}{n_s}$	
<b>HT</b> $\frac{\text{number of turns in primary coil}}{\text{number of turns in secondary coil}}$	<b>HT</b> $\frac{\text{potential difference across primary coil} \times \text{current in primary coil}}{\text{coil} \times \text{current in secondary coil}} = V_p I_p = V_s I_s$



**BLANK PAGE**

**Physics Equations Sheet - GCSE Physics (8463)  
FOR USE IN JUNE 2024 ONLY**

**WP/M/CH/Jun24/8463/INS/V2**