Version 1.0: 0106

ASSESSMENT and QUALIFICATIONS ALLIANCE

General Certificate of Education

Statistics 6380

SS03 Statistics 3

Mark Scheme

2006 examination – January series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Copyright © 2006 AQA and its licensors. All rights reserved.

Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation

√or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
–*x* EE	deduct *x* marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

SS03

Q	Solution	Marks	Total	Comments
1(a)	r = – 0.708 (3 sf from calc) or r = $\dfrac{3527.4-(\frac{115\times 348.1}{11})}{\sqrt{804.727}\times\sqrt{30.96}}$ = $\dfrac{-111.827}{\sqrt{804.727}\times\sqrt{30.96}}$ = – 0.708	B3 or B1 M1 A1	**3**	Alternative $n = 11$ $\sum y = 348.1$ $\sum x = 115$ $\sum y^2 = 11046.75$ $\sum x^2 = 2007$ $\sum xy = 3527.4$ B1 – 0.71 SC B1M1A0 0.708 SC B1M1A0
(b)	H_o $\rho = 0$ H_1 $\rho < 0$ 1 tail 1 % sig level test stat r = - 0.708 cv = – 0. 6851 since ts < – 0.6851 Reject H_o Significant evidence at 1% level to suggest a negative linear association between the age at which a baby first learns to crawl and the average daily temperature during the sixth month of its life.	B1 B1 M1 A1 E1	**5**	for cv for comparison ts/cv not +cv / – pmcc in context EO if x / y used
(c)	A Type I error occurs when the Null Hypothesis is incorrectly rejected: in this case, when the conclusion made is that there is a negative association between temperature and age but, in fact, a negative association does not exist.	E1 E1	**2**	in context Condone x / y Allow 2 tail conclusion
	Total		**10**	

SS03 (cont)

Q	Solution	Marks	Total	Comments
2(a)	H_o pop median difference, $\eta_d = 0$ H_1 pop median difference, $\eta_d \neq 0$ 2 tail 5%	B1		
	Tyre 1 2 3 4 5 6 diff 10.2 2.3 - 0.6 4.3 - 0.8 2.2 rank 12 7½ - 1 9 - 2 6	M1		For differences
	7 8 9 10 11 12 5 - 2.3 - 1.2 8.4 2.1 2.0 10 - 7½ - 3 11 5 4	m1 M1		For ranks of differences For ties
	$T_+ = 12 + 7½ + \ldots + 5 = 64½$ $T_- = 1 + 2 + 7½ + 3 = 13½$	m1		For totals
	Test stat T = 13½	A1		For one correct total
	cv = 14	B1		For cv
	T < 14 Reject H_o	M1		Comparison cv/ts
	Significant evidence at 5% level to suggest that there is a difference in average treadwear measurement for the two methods.	A1 E1	**10**	In context
(b)	In the original design, the same tyre is used each time which eliminates any individual differences between tyres and means that any difference due to measurement method is more likely to be detected, if one exists.	E1		
	If two separate tyres were used, even from the same car and both from the front of the car, there may well be individual differences between them.	E1	**2**	
(c)	Max T = $\sum_{r=1}^{12} r = 1 + 2 + \ldots + 12 = 78$	M1 A1	**2**	
	Total		**14**	

SS03 (cont)

Q	Solution	Marks	Total	Comments
3(a)	The frequencies are very low in several categories (insufficient data) and so a lot of pooling might be necessary that could reduce the contingency table below the 2×2 minimum required to sensibly carry out such an analysis.	B1		All E_i are below 5 and pooling will not solve this problem
	or	E1		any two valid reasons with explanation of reason in context
	The level of poultry in the meat hot dogs is variable – could be 0% or up to 25% - so conclusion would not be relevant to investigating link to actual amount of poultry and sodium levels.	B1		
	or			
	The sodium level categories are not discrete so some hot dogs could have been 'double counted'.	E1	**4**	
(b)	H_o Samples are taken from identical populations			
	H_1 Samples are not taken from identical populations – population average calorie content is lower for poultry hot dogs sausages.	B1		Hypotheses referring to population averages also acceptable
	1 tail 5%	B1		1 tail / ok generous
	Ranks Beef 15 13 12 8 14 16 4 7 Poultry 6 9 2 3 1 10 5 11	M1M1		For ranks as one group – at least 10 correct Other alternative methods acceptable
	$T_B = 15 + 13 + \ldots + 7 = 89$ $T_P = 6 + 9 + \ldots + 11 = 47$	m1		For totals of ranks in each group
	$U_B = 89 - \frac{8\times9}{2} = 53$ $U_P = 47 - \frac{8\times9}{2} = 11$	m1		For U attempted
	Test stat U = 11	A1		For U correct – either
	Cv = 16	B1		For consistent cv with U
	U < 16	M1		For comparison U/cv
	Reject H_o	A1		
	Significant evidence at the 5% level to suggest that the population average calorie content for poultry hot dogs is lower than that for beef hot dogs.	E1	**11**	In context
			15	

SS03 (cont)

Q	Solution	Marks	Total	Comments
4(a)	H_0 Samples from identical populations H_1 Samples not from identical populations 5% sig level	B1 B1		or H_0 $\eta_A = \eta_B = \eta_C$ H_1 at least two of η_A, η_B, η_C do differ Allow $\eta_A \neq \eta_B \neq \eta_c$
	Ranks **Fish Market A** / **Fish Market B** / **Fish Market C** 3 / 1 / 10 6 / 2 / 11 8 / 4 / 13 9 / 5 / 14 12 / 7 / 15	M1 M1		For ranks all as one group – can be reversed For at least 8 correct CAO
	$T_A = 38$ $T_B = 19$ $T_C = 63$ $n_A = 5$ $n_B = 5$ $n_C = 5$	m1 A1		totals any one correct
	$\sum_{i=1}^{m} \frac{T_i^2}{n_i} = \frac{38^2}{5} + \frac{19^2}{5} + \frac{63^2}{5} = 1154.8$	m1		
	$H = \frac{12}{15 \times 16} \times 1154.8 - (3 \times 16)$ $= 9.74$	m1 A1		9.60 – 9.80
	Critical value from $\chi^2_2 = 5.99$ $H > 5.99$	B1 M1		
	Sig evidence to reject H_0 and conclude that samples are not from identical populations	A1		
	There is significant evidence that at least two of the average prices (from Fish Markets A, B or C) do differ.	E1		Difference in context Mention of 'at least two' √ E1, E0 if Accept H_0
		E1	**14**	Significant evidence to suggest that the mean price for C is certainly greater than the mean price for B
(b)	Medians 227.3 , 223.4, 249.6 It would appear that average prices at Fish Market C were significantly higher (as there is significant evidence of a difference detected in part (a)) and this would be the recommended Fish Market for Chinook salmon	B1 E1	**2**	Identification of C with reason – generous
			16	

SS03 (cont)

Q	Solution	Marks	Total	Comments
5(a)(i)	H_o Violence level is independent of type of offence H_1 Violence level is not independent of type of offence 1 tail 5%	B1		
		M1		M1 E method for 5 correct
		m1		For all E correct
	Two E_i in the 'violence involving weapons' column are below 5 so pooling is required			
		m1		For pooling
		A1		
	$ts = \sum \frac{(O-E)^2}{E}$ $= \frac{2.53^2}{55.47} + \frac{2.53^2}{19.53} + \frac{4.34^2}{47.34} + \frac{4.34^2}{16.66}$ $\frac{1.81^2}{22.19} + \frac{1.81^2}{7.81}$	m1		ts sum with correct denominators (condone no pooling)
	= 2.54	A1		For ts in range 2.30 – 2.80 (or 6.10 - 6.50 ft)
	cv df = 2 5% cv = 5.991	B1		For cv (9.488 ft)
	ts < 5.991 Accept H_o	m1		For comparison ts/cv ft
	No significant evidence to suggest use of violence is associated with type of offence	A1	**10**	sc If ts only sc (m1, 1, m1, A1) 3 6.1 – 6.5 sc (m1, 1, 1 A1, M1, A1) 4 2.3 – 2.8

	No violence	Violence but no weapons	Violence involving weapons
Non drug rel theft or damage	55.47	14.20	5.33
Drug rel theft or damage	47.34	12.12	4.54
Other	22.19	5.68	2.13

	No Violence	Violence
Non drug rel theft or damage	55.47	19.53
Drug rel theft or damage	47.34	16.66
Other	22.19	7.81

SS03 (cont)

Q	Solution	Marks	Total	Comments
(b)(i)	H_o Type of sentence is independent of whether firearms were used H_1 Type of sentence is not independent of whether firearms were used 1 tail 1%	B1		
		B1		For E values method
	$ts = \sum \frac{(\lvert O-E\rvert - 0.5)^2}{E} =$ $\frac{5.5^2}{26} + \frac{5.5^2}{8} + \frac{5.5^2}{52} + \frac{5.5^2}{16} = 7.42$	M1 M1 A1		For ts for Yates' corr For ts 7.2 – 7.7
	cv df = 1 1% cv = 6.635 ts > 6.635	B1 m1		For cv For comparison ts/cv
	Reject H_o Significant evidence to suggest that type of sentence is not independent of whether firearms were used	A1	**8**	
(b)(ii)	Offences where firearms are used are much **more likely** to result in a custodial sentence (and those where firearms are not used are **less likely** to result in a custodial sentence.)	B1 E1	**2**	Correct association identified Explained in context
			20	

	Not used	Used
Non custodial	26	8
Custodial	52	16