



General Certificate of Education (A-level)
January 2013

Statistics

SS04

(Specification 6380)

Statistics 4

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
–x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Marks	Total	Comments
1 (a)	No. turning up (X) ~ B(60, 0.8) which is approximately N(48, 9.6)	B1 M1 A1	6	May be implied Attempt at any normal approximation Mean = 48 cao, variance = 9.6 cao (or SD = 3.09 ~ 3.1(0))
	$P(X > 50) = P(Z > \frac{50.5 - 48}{\sqrt{9.6}})$ $= P(Z > 0.81)$ $= 1 - 0.79103$ $= 0.20897$	M1 m1 A1		Standardizing with their μ and σ (allow missing CC or 49.5); ignore sign Use of 50.5 and correct sign (0.81 not required) AWFW 0.208~0.212 (more exact value 0.2099) (Notes: (i) Use of B(60, 0.8) gives answer 0.21321 scoring 1/6) (ii) Use of 50 or 49.5 gives 0.259 or 0.314 respectively scoring 4/6
(b)	Fairly high probability (of disgruntled customers.)	B1	2	Sensible comment on size of prob (may say 0.21 is 'high' or 'low'; if prob >> 0.21 should be 'high; if prob << 0.21 should be 'low') or using original numbers + some attempt at a conclusion
	No, would not recommend continuing.	Bdep1		All correct (as bracket above) with clear, sensible conclusion (eg fairly low prob so recommend continuing gets 2/2)
Total			8	

Q	Solution	Marks	Total	Comments
2 (a)(i)	Use of 2.5758	B1	4	Accept 2.57~2.58 (even if called <i>t</i>)
	99% CI is $6.75 \pm 2.5758 \times \frac{1.29}{\sqrt{1403}}$	M1		Use of $\frac{1.29}{\sqrt{1403}}$ (= 0.0344)
	= $6.75 \pm 0.088(7)$	m1		Correct interval, allow any Z
	= (6.66, 6.84)	A1		Either for $6.75 \pm (0.088 \sim 0.089)$ or AWR 6.66 and 6.84
	(ii) Comparing upper limit of CI for males with lower limit of CI for females.	M1	2	Comparing limits of two CIs
	There is a difference between males and females.	A1		Requires implication of “difference” and clearly implied comparison of their upper limit with 6.87
	(b)(i) $H_0 : p = 0.15$	B1	6	For both
	$H_1 : p > 0.15$	B1		Use of correct distribution
	Under H_0 , number sleeping over 8 hours ~ Bin(14, 0.15)	M1		For finding $p(X \geq 4)$ or $p(X > 4)$ from a binomial distribution
	Then $p(X \geq 4) = 1 - 0.8535$	A1		(0.146 ~ 0.147)
	= 0.1465	M1		Their binomial prob compared with 0.05 and correct ft conclusion
	Cannot reject H_0 at the 5% level	E1		Correct conclusion in context. Needs all previous marks.
	No evidence of difference from 15 percent.			<i>Note</i> $p(X > 4) = 1 - 0.9533 = 0.047$ so reject H_0 gets B1B1M1A0M1E0 max 4/6
	(ii) Roughly the same age (not all teenagers)	B1	2	Any sensible comment on the restricted nature of Rowan’s sample.
	Same school/location. (not all UK)			
	Similar social backgrounds etc			
	May not know/remember how long they slept.	B1		Any sensible comment on issues such as inability to remember or truthfulness or accuracy.
	May not be truthful.			
Total			14	

Q	Solution	Marks	Total	Comments
3 (a)	Total weight when Safeerah's riding is: 65 + 15 = 80kg on steel bike 65 + 10 = 75 kg on carbon bike Then % reduction = $5/80 \times 100\%$ = 6.25%.	M1 A1	2	For attempt to include Safeerah's weight Correct calculation (Accept "approx 6"). Alternatively, something like "94% of 80 is approx 75"
	Hence Josh's figure			
	(b) $\bar{x} = 88.175$ $s = 4.6809$	B1		For 88.2 and $s_{n-1} = 4.68$ or $s_n = 4.48(16)$ (ignore labels)
	$H_0 : \mu = 90$	B1		Both
	$H_1 : \mu < 90$			
	$t = \frac{88.175 - 90}{4.6809/\sqrt{12}}$	M1		M1 for use of $\frac{s_{n-1}}{\sqrt{n}}$ or $\frac{s_n}{\sqrt{n-1}}$ in test statistic formula
		m1		Correct formula, ignore sign for m1.
				Or $t = \frac{88.175 - 90}{4.4816/\sqrt{11}}$
	= -1.351	A1		AWFW -1.36 to -1.33
	Critical value $t_{11} = -1.796$	B1 B1		For 11 df may be implied (eg by 1.80 or 2.20(1)) For $t_{11} = (-1.80 \sim -1.79)$ or $p = 0.10197$ (AWFW 0.1005 to 0.1055)
	Accept H_0 at 5% level. No evidence that mean journey time is < 90 minutes.	E1	8	Their t and critical t (both negative) OR their p-value and 0.05 but still requires negative t.. Requires M1 and m1 and "acc H_0 ". In context Alternatives Full marks available for: (i) one-sided CI $88.175 + 2.427 = 90.6$ then cf 90 (ii) critical region $90 - 2.427 = 87.6$ then cf 88.175
	Total		10	

Q	Solution	Marks	Total	Comments
4	(a) Distributed at random/independently/ at constant average rate.	B1	1	For any of these.
	(b) $H_0 : \lambda = 12$ $H_1 : \lambda < 12$ Find $P(X \leq 8)$ from Poisson tables $= 0.155(0)$ This is > 0.10 so no evidence it's reduced.	B1 M1 M1 A1	4	For both. Allow μ or "rate". Attempt to calculate $P(X \leq 8)$ or $P(X < 8)$ ($= 0.0895$) Compare their Poisson prob with 0.10 All correct including 0.155. Allow "accept H_0 " here and isw.
	(c) Number of lob worms in 10 m^2 (Y) has a Poisson distribution with $\lambda = 142$ Then 95% CI for Y using $Z=1.96$ is $142 \pm 1.96\sqrt{142}$ $= 142 \pm 23.356$ Then average density per m^2 is 14.2 ± 2.34 (11.86, 16.54)	B1 M1 A1 M1 A1	5	For 1.96 For $c \pm z\sqrt{c}$. Allow $c \pm z\sqrt{\frac{c}{10}}$ $142 \pm (23.3 \sim 23.4)$ or (119, 165) Dividing this (or limits) by 10 For AWRT (11.9, 16.5) or $14.2 \pm (2.3 \sim 2.4)$ Notes: (i) SR: Using $14.2 \pm 1.96\sqrt{14.2}$ gets B1M1A0 then B1 for answer 14.2 ± 7.4 getting max 3/5. (ii) Using $14.2 \pm 1.96\sqrt{\frac{14.2}{10}}$ gets potentially full marks.
	(d) (i) $M_A = 56$ and $M_B = 13$ (ii) Would expect $V_A > M_A$ Ants 'attract' each other \rightarrow tendency for small or large numbers in a sq m \rightarrow larger variation (than expected from Poisson)	B1 B1 B1	5	For both Or equiv. Any indication of why they would expect "large" variation (more than just saying they live in colonies). Note: B1's indep in either order
	(iii) Would expect $V_B < M_B$ Beetles 'repel' each other \rightarrow more evenly spread (eg unlikely to have high concentrations) \rightarrow smaller variation (than expected from Poisson).	B1 B1	5	Or equiv. Any indication of why they would expect "small" variation (more than just saying they are territorial). As note in (ii).
	Total		15	

Q	Solution	Marks	Total	Comments
5 (a)	P(dizziness) \times P(swollen jnts) = $0.12 \times 0.15 = 0.018$ = given value for joint probability. Hence independent.	M1 A1	2	For 0.12×0.15 On correct reasoning.
(b) (i)	No. with dry skin $\sim B(90, 0.01)$ Which is approx Poisson with $\lambda = 90 \times 0.01 = 0.9$ Then $P(X > 2) = 1 - 0.937(1)$ = 0.0629	B1 B1dep M1 A1	4	For Poisson (may be implied) May be implied Allow for $1 - 0.7725$ (may be implied by $0.227 \sim 0.228$) $0.0629 \sim 0.063$ Note: Exact binom $\rightarrow 0.0619 \rightarrow (0/4)$
(ii)	$H_0 : p = 0.12$ $H_1 : p \neq 0.12$ $X \sim B(90, 0.12) \approx N(10.8, 9.504)$ $TS = \frac{21 - 10.8}{\sqrt{9.504}} = 3.31$ or $TS = \frac{20.5 - 10.8}{\sqrt{9.504}} = 3.15$ or using proportions Use of $N(0.12, .00117)$ $\hat{p} = \frac{21}{90} = 0.233$ $TS = \frac{0.233 - 0.12}{\sqrt{\frac{0.12 \times 0.88}{90}}} = 3.31$ Critical values are ± 2.5758 Reject H_0 at the 1% level There is evidence that the proportion of patients really does differ from 0.12.	B1 B1 B1 M1 A1 (M1) (A1) (B1) (B1) (M1) (A1) B1 A1 E1	8	Both B1 for mean 10.8 cao B1 for variance ($9.5(0) \sim 9.51$) or SD ($3.08 \sim 3.09$) Their mean/SD $3.25 \sim 3.4$ $3.13 \sim 3.15$ Note: wrong CC (21.5) gives 3.47 for M1A0 B1 for mean 0.12 cao B1 for variance ($0.0011 \sim 0.0012$) or SD ($0.033 \sim 0.035$) Their mean/SD $3.25 \sim 3.4$ $2.57 \sim 2.58$ or $p = 0.000938$ (AWFW 0.00065 to 0.00115) from $TS = 3.31$ or $p = 0.00163$ (AWFW 0.00163 to 0.00175) from $TS = 3.15$ Allow $p = 0.000466$ or 0.000816 only if compared with 0.005 fit their TS and critical z value or their p -value and 0.01 or their $\frac{1}{2} p$ and 0.005. Must correctly reject H_0 In context, requires all previous marks in (b)(ii).
	Total		14	

Q	Solution	Marks	Total	Comments
6(a) (i)	Let L be the height of cut trees Then $L = H + G - S$ $E(L) = 50 + 150 - 10 = 190$	B1	3	For 190 cao. Method must be seen and correct. Correct expression for V(L) seen AWRT 3.10 answer given.
	$V(L) = 1.49^2 + 2.67^2 + 0.51^2 (= 9.6091)$ Then SD(L) = 3.10 as required.	M1 A1		
(ii)	$P(L < 1.95) = P(Z < \frac{195 - 190}{3.10})$	M1	2	Allow wrong tail but needs 195, 190 and 3.10 or their E(L)&SD(L) (0.946 ~ 0.947) SC (1.95-1.9)/3.1 = 0.0161 gets B1
	$= P(Z < 1.61(3))$ $= 0.9463(0)$	A1		
(b) (i)	$P(97 < \text{height} < 103) = P(-1 < Z < 1)$		3	Correct method for z and 0.841 seen. (0.682 ~ 0.683) For their probability $\times 100$. May be rounded up or down.
	$= 0.84134 - (1 - 0.84134)$ $= 0.68268$ Can expect $100 \times 0.683 = 68.3$ trees	M1 A1 B1		
(ii)	Let P be the tree length for shredding Then $E(P) = 190 - 100 = 90$ and $V(P) = 3.1^2 + 3^2 = 18.61$		4	Marks in (ii) can be earned in (iii) No marks yet! Need to use these values for income. For 500 + attempt at pulp income
	Income (Y) = $500 + 100 \times 0.02 \times P$ $= 500 + 2P$ Then $E(Y) = 500 + 2 \times 90 = 680$ and $V(Y) = 2^2 \times V(P) = 74.44$	M1 A1 AF1		
(iii)	Thus $Y \sim N(680, 74.44)$	B1	2	For normal distribution Their E(Y) compared with 750 (may be implied) All clear & correct, including 680 Alternative May use their distribution to find a probability (eg $P(Y > 750) \approx 0$). Then M1 for attempt at relevant probability and E1 as above. SC Something like “no guarantee she will sell them all so should take what is firmly offered” B1
	Since council's offer gets $\pounds 680 < \pounds 750$ she was NOT wise to take it up.	M1 E1		
	Total		14	
	TOTAL		75	