

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
TOTAL	



General Certificate of Education
Advanced Level Examination
June 2010

Mathematics

MFP2

Unit Further Pure 2

Wednesday 9 June 2010 1.30 pm to 3.00 pm

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.



J U N 1 0 M F P 2 0 1

Answer **all** questions in the spaces provided.

1 (a) Show that

$$9 \sinh x - \cosh x = 4e^x - 5e^{-x} \quad (2 \text{ marks})$$

(b) Given that

$$9 \sinh x - \cosh x = 8$$

find the exact value of $\tanh x$. (7 marks)

<small>QUESTION PART REFERENCE</small>	



QUESTION
PART
REFERENCE

This section contains a large grid for writing answers. The grid consists of 28 rows. The first column is a narrow vertical column for writing the question reference, and the remaining 27 columns are wide enough for a full answer. Each row is bounded by a solid horizontal line at the top and a solid horizontal line at the bottom, with a dotted horizontal line in the middle of each row. The entire grid is enclosed in a rectangular border.

Turn over ►



3 Two loci, L_1 and L_2 , in an Argand diagram are given by

$$L_1 : |z + 1 + 3i| = |z - 5 - 7i|$$

$$L_2 : \arg z = \frac{\pi}{4}$$

(a) Verify that the point represented by the complex number $2 + 2i$ is a point of intersection of L_1 and L_2 . (2 marks)

(b) Sketch L_1 and L_2 on one Argand diagram. (5 marks)

(c) Shade on your Argand diagram the region satisfying

both $|z + 1 + 3i| \leq |z - 5 - 7i|$

and $\frac{\pi}{4} \leq \arg z \leq \frac{\pi}{2}$ (2 marks)

QUESTION
PART
REFERENCE



4 The roots of the cubic equation

$$z^3 - 2z^2 + pz + 10 = 0$$

are α , β and γ .

It is given that $\alpha^3 + \beta^3 + \gamma^3 = -4$.

(a) Write down the value of $\alpha + \beta + \gamma$. *(1 mark)*

(b) (i) Explain why $\alpha^3 - 2\alpha^2 + p\alpha + 10 = 0$. *(1 mark)*

(ii) Hence show that

$$\alpha^2 + \beta^2 + \gamma^2 = p + 13$$
 (4 marks)

(iii) Deduce that $p = -3$. *(2 marks)*

(c) (i) Find the real root α of the cubic equation $z^3 - 2z^2 - 3z + 10 = 0$. *(2 marks)*

(ii) Find the values of β and γ . *(3 marks)*

QUESTION
PART
REFERENCE



QUESTION
PART
REFERENCE

A large rectangular area with a vertical line on the left side and horizontal dotted lines across the page, intended for writing answers.



Turn over ►

5 (a) Using the identities

$$\cosh^2 t - \sinh^2 t = 1, \quad \tanh t = \frac{\sinh t}{\cosh t} \quad \text{and} \quad \operatorname{sech} t = \frac{1}{\cosh t}$$

show that:

(i) $\tanh^2 t + \operatorname{sech}^2 t = 1;$ *(2 marks)*

(ii) $\frac{d}{dt}(\tanh t) = \operatorname{sech}^2 t;$ *(3 marks)*

(iii) $\frac{d}{dt}(\operatorname{sech} t) = -\operatorname{sech} t \tanh t.$ *(3 marks)*

(b) A curve C is given parametrically by

$$x = \operatorname{sech} t, \quad y = 4 - \tanh t$$

(i) Show that the arc length, s , of C between the points where $t = 0$ and $t = \frac{1}{2} \ln 3$ is given by

$$s = \int_0^{\frac{1}{2} \ln 3} \operatorname{sech} t \, dt$$
 (4 marks)

(ii) Using the substitution $u = e^t$, find the exact value of s . *(6 marks)*

QUESTION
PART
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6 (a) Show that $\frac{1}{(k+2)!} - \frac{k+1}{(k+3)!} = \frac{2}{(k+3)!}$. (2 marks)

(b) Prove by induction that, for all positive integers n ,

$$\sum_{r=1}^n \frac{r \times 2^r}{(r+2)!} = 1 - \frac{2^{n+1}}{(n+2)!} \quad (6 \text{ marks})$$

QUESTION
PART
REFERENCE

Area with horizontal dotted lines for writing answers.



7 (a) (i) Express each of the numbers $1 + \sqrt{3}i$ and $1 - i$ in the form $re^{i\theta}$, where $r > 0$.
(3 marks)

(ii) Hence express

$$(1 + \sqrt{3}i)^8(1 - i)^5$$

in the form $re^{i\theta}$, where $r > 0$. (3 marks)

(b) Solve the equation

$$z^3 = (1 + \sqrt{3}i)^8(1 - i)^5$$

giving your answers in the form $a\sqrt{2}e^{i\theta}$, where a is a positive integer and
 $-\pi < \theta \leq \pi$. (4 marks)

QUESTION
PART
REFERENCE



QUESTION
PART
REFERENCE

A large rectangular area with horizontal dotted lines for writing, intended for answers to questions.

END OF QUESTIONS



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

