



**General Certificate of Education (A-level)  
June 2011**

**Mathematics**

**MFP2**

**(Specification 6360)**

**Further Pure 2**

**Final**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2011 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

MFP2

Q	Solution	Marks	Total	Comments
1(a)				Use average of whole question if 2 diagrams used
(i)	Circle correct centre touching $x$ -axis	B1 B1 B1F	3	Circle in any position Must be shown ft incorrect centre
(ii)	half-line through $(0, -2)$ through point of contact of circle with $x$ -axis	B1 B1 B1	3	Can be inferred
(b)	Inside circle On line	B1 B1F	2	ft errors in position of line and circle
<b>Total</b>			<b>8</b>	
2(a)	$\frac{(e^x + e^{-x})}{2} \frac{(e^y + e^{-y})}{2} - \frac{(e^x - e^{-x})}{2} \frac{(e^y - e^{-y})}{2}$ <p>Correct expansions</p> $= \frac{1}{2}(e^{x-y} + e^{-(x-y)}) = \cosh(x-y)$	M1A1 A1 A1	4	M0 if sinh and cosh confused M1 for formula quoted correctly Use of $e^{xy}$ A0 AG
(b)(i)	$\cosh(x - \ln 2) = \cosh x \cosh(\ln 2) - \sinh x \sinh(\ln 2)$ $\left. \begin{aligned} \cosh(\ln 2) &= \frac{5}{4} \\ \sinh(\ln 2) &= \frac{3}{4} \end{aligned} \right\} \text{any method}$ $\frac{5}{4} \cosh x = \frac{7}{4} \sinh x$ $\tanh x = \frac{5}{7}$	M1 B1 A1F A1	4	<p><b>Alternative:</b></p> $\frac{e^{x-\ln 2} + e^{-x+\ln 2}}{2} = \frac{e^x - e^{-x}}{2} \quad \text{M1}$ <p>Both correct  <math display="block">e^{x-\ln 2} = \frac{e^x}{2} \text{ or } e^{-x+\ln 2} = 2e^{-x}</math> used B1</p> $e^x = \sqrt{6} \quad \text{A1}$ $\tanh x = \frac{5}{7} \quad \text{A1}$
(ii)	$x = \frac{1}{2} \ln \left( \frac{1 + \frac{5}{7}}{1 - \frac{5}{7}} \right) \text{ or } \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{5}{7}$ $= \frac{1}{2} \ln 6$	M1 A1	2	Could be embedded in (b)(i)
<b>Total</b>			<b>10</b>	

**MFP2 (cont)**

Q	Solution	Marks	Total	Comments	
<b>3(a)</b>	$(r+1)! = (r+1)r(r-1)!$	M1	2	AG	
	Result	A1			
	<b>(b)</b> <b>Attempt</b> to use method of differences	$\sum_{r=1}^n (r^2 + r - 1)(r-1)! = (n+1)! + n! - 1! - 0!$	M1	4	Must be seen AG
		$(n+1)! = (n+1)n!$	m1		
		$(n+2)n! - 2$	A1		
<b>Total</b>			<b>6</b>		
<b>4(a)(i)</b>	$\sum \alpha = 2$	B1	2		
	$\sum \alpha\beta = 0$	B1			
<b>(ii)</b>	$\sum \alpha^2 = (\sum \alpha)^2 - 2\sum \alpha\beta$	M1	2	Used. Watch $\sum \alpha = -2$ (M1A0) AG	
	$= 4$	A1			
<b>(iii)</b>	Clear explanation	E1	1	eg $\alpha$ satisfies the cubic equation since it is a root. Accept $z = \alpha$	
<b>(iv)</b>	$\sum \alpha^3 = 2\sum \alpha^2 - 3k$	M1	2	Or $\sum \alpha^3 = (\sum \alpha)^3 - 3\sum \alpha \sum \alpha\beta + 3\alpha\beta\gamma$ AG	
	$= 8 - 3k$	A1			
<b>(b)(i)</b>	$\alpha^4 = 2\alpha^3 - k\alpha$	B1	4	Or $\sum \alpha^4 = (\sum \alpha^2)^2 - 2(\sum \alpha\beta)^2 + 4\alpha\beta\gamma \sum \alpha$ ft on $\sum \alpha = -2$ AG	
	$\sum \alpha^4 = 2\sum \alpha^3 - k\sum \alpha$	M1			
	$= 2(8 - 3k) - 2k$	A1			
	$k = 2$	A1			
<b>(ii)</b>	$\sum \alpha^5 = 2\sum \alpha^4 - k\sum \alpha^2$	M1	3		
	Substitution of values	A1			
	$= -8$	A1			
<b>Total</b>			<b>14</b>		

MFP2 (cont)

Q	Solution	Marks	Total	Comments
5(a)	$2y \frac{dy}{dx} = 2x$ $S = 2\pi \int_0^6 y \sqrt{1 + \frac{x^2}{y^2}} dx$	B1 M1 A1F	5	Or $\frac{dy}{dx} = x(x^2 + 8)^{-\frac{1}{2}}$ M1 for use of formula provided $\frac{dy}{dx}$ is a function of $x$ A1 for substitution for $\frac{dy}{dx}$ (one slip)
	Eliminating all $y$ $= 2\sqrt{2}\pi \int_0^6 \sqrt{x^2 + 4} dx$	m1 A1		AG
(b)	$dx = 2 \cosh \theta d\theta$ or $\frac{dx}{d\theta} = 2 \cosh \theta$	B1	8	For eliminating $x$ completely and use of $d\theta$ , ie $d\theta$ attempted Use of $\cosh^2 \theta - \sinh^2 \theta = 1$ (ignore limits) Use of formula for $\cosh 2\theta$ ; must be correct Correct integration of $a \cosh 2\theta + b$ Use of $\sinh 2\theta = 2 \sinh \theta \cosh \theta$ Must be seen Or change limits
	$S = 2\sqrt{2}\pi \int \sqrt{4\sinh^2 x + 4} \cdot 2\cosh \theta d\theta$	M1		
	$S = (2\sqrt{2})\pi \int 2\cosh \theta \cdot 2\cosh \theta d\theta$	m1		
	$= 4\sqrt{2}\pi \int (\cosh 2\theta + 1) d\theta$	m1		
	$= 4\sqrt{2}\pi \left[ \frac{\sinh 2\theta}{2} + \theta \right]$	B1F		
	$= 4\sqrt{2}\pi [\sinh \theta \cosh \theta + \theta]$	m1		
$= 4\sqrt{2}\pi \left[ \frac{x}{2} \sqrt{\frac{x^2}{4} + 1} + \sinh^{-1} \frac{x}{2} \right]_0^6$	M1			
$= \pi [24\sqrt{5} + 4\sqrt{2} \sinh^{-1} 3]$	A1	AG		
<b>Total</b>			<b>13</b>	
6(a)	Expansion of $(k+1)(4(k+1)^2 - 1)$ $= 4k^3 + 12k^2 + 11k + 3$	M1 A1	2	Any valid method – first step correct AG
	Assume true for $n=k$ For $n=k+1$ : $\sum_{r=1}^{k+1} (2r-1)^2 = \frac{1}{3}k(4k^2 - 1) + (2k+1)^2$ $= \frac{1}{3}(4k^3 + 12k^2 + 11k + 3)$ $= \frac{1}{3}(k+1)(4(k+1)^2 - 1)$	M1A1 A1F		
(b)	True for $n=1$ shown Proof by induction set out properly (if factorised by 3 linear factors, allow A1 at this particular point)	B1 E1	6	Using part (a) Dependent on all marks correct
<b>Total</b>			<b>8</b>	

**MFP2 (cont)**

Q	Solution	Marks	Total	Comments
7(a)(i)	$\cos 5\theta + i \sin 5\theta = (\cos \theta + i \sin \theta)^5$	M1	5	Attempt to expand 3 correct terms
	Expansion in any form Equate real parts: $\cos 5\theta = \cos^5 \theta - 10 \cos^3 \theta \sin^2 \theta + 5 \cos \theta \sin^4 \theta$	A1 m1 A1		Correct simplification AG
(ii)	Equate imaginary parts: $\sin 5\theta = 5 \cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin^5 \theta$	A1	3	CAO
	$\tan 5\theta = \frac{\sin 5\theta}{\cos 5\theta}$ Division by $\cos^5 \theta$ or by $\cos^4 \theta$ $\tan 5\theta = \frac{\tan \theta (5 - 10 \tan^2 \theta + \tan^4 \theta)}{1 - 10 \tan^2 \theta + 5 \tan^4 \theta}$	M1 m1 A1	3	Used AG
(b)	$\theta = \frac{\pi}{5} \Rightarrow \tan 5\theta = 0$	M1	3	Or for $\tan^4 \theta - 10 \tan^2 \theta + 5 = 0$
	$\therefore \tan \frac{\pi}{5}$ satisfies $t^4 - 10t^2 + 5 = 0$	A1		Or for $\tan 5\theta = 0$
(c)	Other roots $\tan \frac{k\pi}{5} \quad k=2, 3, 4$	B1	3	OE
	Product of roots = 5	M1	5	Or $\tan \frac{2\pi}{5} = -\tan \frac{3\pi}{5}$
	$\tan \frac{\pi}{5} = -\tan \frac{4\pi}{5}$	B1		
	$\tan^2 \frac{\pi}{5} \tan^2 \frac{2\pi}{5} = 5$	A1		
	$\tan \frac{\pi}{5} \tan \frac{2\pi}{5} = +\sqrt{5}$	A1		
- sign rejected with reason	E1			
	<b>Alternative (c)</b> Use of quadratic formula $t^2 = 5 \pm 2\sqrt{5}$ $t = \pm\sqrt{5 \pm 2\sqrt{5}}$ Correct selection of +ve values Multiplied together to get $\sqrt{5}$	M1 A1 B1 E1 A1		
	<b>Total</b>		<b>16</b>	
	<b>TOTAL</b>		<b>75</b>	