

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



General Certificate of Education
Advanced Subsidiary Examination
June 2011

Mathematics

MPC1

Unit Pure Core 1

Wednesday 18 May 2011 9.00 am to 10.30 am

<p>For this paper you must have:</p> <ul style="list-style-type: none"> the blue AQA booklet of formulae and statistical tables. <p>You must not use a calculator.</p>	
---	--

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.
- The use of calculators is **not** permitted.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.



J U N 1 1 M P C 1 0 1

- 5** The polynomial $p(x)$ is given by $p(x) = x^3 - 2x^2 + 3$.
- (a)** Use the Remainder Theorem to find the remainder when $p(x)$ is divided by $x - 3$. *(2 marks)*
- (b)** Use the Factor Theorem to show that $x + 1$ is a factor of $p(x)$. *(2 marks)*
- (c) (i)** Express $p(x) = x^3 - 2x^2 + 3$ in the form $(x + 1)(x^2 + bx + c)$, where b and c are integers. *(2 marks)*
- (ii)** Hence show that the equation $p(x) = 0$ has exactly one real root. *(2 marks)*

QUESTION
PART
REFERENCE



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

