



**General Certificate of Education (A-level)
June 2011**

Mathematics

MS2B

(Specification 6360)

Statistics 2B

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

| | |
|--------------|--|
| M | mark is for method |
| m or dM | mark is dependent on one or more M marks and is for method |
| A | mark is dependent on M or m marks and is for accuracy |
| B | mark is independent of M or m marks and is for method and accuracy |
| E | mark is for explanation |
| ✓ or ft or F | follow through from previous incorrect result |
| CAO | correct answer only |
| CSO | correct solution only |
| AWFW | anything which falls within |
| AWRT | anything which rounds to |
| ACF | any correct form |
| AG | answer given |
| SC | special case |
| OE | or equivalent |
| A2,1 | 2 or 1 (or 0) accuracy marks |
| -x EE | deduct x marks for each error |
| NMS | no method shown |
| PI | possibly implied |
| SCA | substantially correct approach |
| c | candidate |
| sf | significant figure(s) |
| dp | decimal place(s) |

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MS2B

| Q | Solution | Marks | Total | Comments |
|---------|--|--------------------------|-----------|--|
| 1(a)(i) | $X \sim \text{Po}(13)$ | B1 | 1 | Both Poisson and $\lambda = 13$ |
| (ii) | $P(X = 20) = P(X \leq 20) - P(X \leq 19)$ $= 0.975(0) - 0.957(3)$ [allow 0.975 - 0.957] $= 0.0177$ (3sf) | M1 A1 | 2 | Must use $\lambda = 13$ otherwise M0A0 AWFW 0.0176 to 0.018 or $P(X = 20) = \frac{e^{-13} \times 13^{20}}{20!}$ M1 $= 0.0177$ A1 |
| (iii) | $P(6 \leq X \leq 18) = P(X \leq 18) - P(X \leq 5)$ $= 0.930(2)$ $- (0.0107 \text{ or } 0.0259)$ $= 0.920$ (3sf) | M1 M1 A1 | 3 | AWFW 0.919 to 0.92 |
| (b) | Cars not random Cars not independent Mean and Variance of cars different / not equal Mean / Average / λ / 2.6 greater / less / smaller / different / variable / not constant / too small / too large Any contextual reason that suggests a change in traffic flow, eg due to: rush hour / congestion / traffic jams / accidents / work traffic / school traffic / peak time | B1 B1 | 2 | Allow (number of) cars not random / not independent B1 for any one of these 3 statements Must indicate a reference to cars Correct comment about value of $\lambda \neq 2.6$ Any combination (one from each group): eg mean greater <i>due to</i> rush hour, or λ smaller <i>due to</i> congestion, or 2.6 too small <i>due to</i> school traffic |
| (c) | $Y \sim \text{Bin}(20, 0.2)$ $P(Y \geq 5) = 1 - P(Y \leq 4)$ $= 1 - 0.6296$ $= 0.37(0)$ (3sf) | M1 A1 | 2 | or: $1 - \left(\begin{array}{l} 0.01153 + 0.05765 + 0.13691 \\ + 0.20536 + 0.21820 \end{array} \right)$ $1 - 0.6296$ (Allow $1 - 0.8042$ seen for M1) |
| (d) | X and Y independent $p = 0.0177 \times 0.3704$ $= 0.00656$ (3sf) | B1 M1 A1 | 3 | Any statement which indicates two / both events are independent [their (c)] \times [their (a)(ii)] AWFW 0.0065 and 0.0067 |
| | | | 13 | |

MS2B (cont)

| Q | Solution | Marks | Total | Comments |
|--------------|--|----------------|-----------|---|
| 2(a)(i) | Area / F(x) = 10u × 0.01π (OE) = 1 ⇒ u = $\frac{10}{\pi}$ | B1 | 2 | Shown by any correct method Alternatives: $f = \frac{1}{10u}$ B1 Show $u = \frac{10}{\pi}$ or show $\frac{1}{10u} = 0.01\pi$ Bdep1 |
| | or $u = \frac{10}{\pi} \Rightarrow F(x) = 1$ | (Bdep1) | | |
| (ii) | $E(X) = \frac{1}{2}(11u + u) = 6u = 6 \times \frac{10}{\pi} = \frac{60}{\pi}$ | B1 | 1 | Must be in terms of π (eg 60π ⁻¹) Alternatives: $\frac{10000}{12\pi^2} = \frac{5000}{6\pi^2} = \frac{2500}{3\pi^2} = \left(\frac{50}{\pi\sqrt{3}}\right)^2 = \frac{(AWRT\ 833)}{\pi^2}$ Must be in terms of π |
| | $Var(X) = \frac{1}{12}(b-a)^2$ $Var(X) = \frac{1}{12}(11u-u)^2$ $= \frac{1}{12} \times 100 \times \frac{100}{\pi^2} = \frac{100^2}{12\pi^2}$ | B1 | | |
| (iii) | $C = \pi \left(X + \frac{10}{\pi} \right)$ | | | |
| | $E(C) = \left. \begin{array}{l} \pi \times [\text{their } E(X)] + 10 \\ \pi \times \frac{60}{\pi} + 10 \end{array} \right\}$ | M1 | | Their numerical value of E(X) used correctly Must have a multiplier of π or 2π |
| | = 70 | A1 | | CAO |
| | $Var(C) = \pi^2 \times \frac{100^2}{12\pi^2} = \frac{100^2}{12}$ $= 833\frac{1}{3} \text{ (833.}\dot{3}\text{)}$ | M1 A1 | 4 | $\pi^2 \times [\text{their } Var(X) > 0]$ Must have a multiplier of π ² or 4π ² Alternatives: $\frac{10000}{12} = \frac{5000}{6} = \frac{2500}{3}$ Must be exact: 833.3 gets A0 |
| (b) | n = 100 and $\bar{a} = 40.5$ $95\% \text{ CI for } \mu = \left. \begin{array}{l} 40.5 \pm z \times \frac{\sqrt{25}}{\sqrt{100}} \\ 40.5 \pm 1.0 \end{array} \right\}$ = (39.5, 41.5) | B1 M1 A1 | 3 | For z = 1.96 z = 1.96 or 1.64 to 1.65 only AWRT |
| Total | | | 11 | |

MS2B (cont)

| Q | Solution | Marks | Total | Comments | | | | | | | | | | | | | | | | | | |
|--------------|---|-------------------------|------------------|--|-----------------|---|---|---|----|----|------------------|----|---|---|----------------|----------------|----------------|-----------------|-----------------|----------------|---|--|
| 4(a) | $E(X) = \sum xp$ $= \frac{3}{40} + \left(2 \times \frac{6}{40}\right) + \left(3 \times \frac{9}{40}\right) + \left(4 \times \frac{12}{40}\right) + \left(5 \times \frac{5}{20}\right) = 3.5$ | B2,1 | 2 | | | | | | | | | | | | | | | | | | | |
| (b)(i) | $E\left(\frac{1}{X}\right) = \sum \frac{1}{x} \times p$ $= \left(1 \times \frac{3}{40}\right) + \left(\frac{1}{2} \times \frac{6}{40}\right) + \left(\frac{1}{3} \times \frac{9}{40}\right) + \left(\frac{1}{4} \times \frac{12}{40}\right) + \left(\frac{1}{5} \times \frac{5}{20}\right)$ $= \frac{7}{20}$ | M1 A1 | 2 | At least 4 of these terms added (accept decimal equivalents) AG (allow 0.35 seen) | | | | | | | | | | | | | | | | | | |
| (ii) | $E\left(\frac{1}{X^2}\right) = \sum \frac{1}{x^2} \times p$ $= \left(1 \times \frac{3}{40}\right) + \left(\frac{1}{4} \times \frac{6}{40}\right) + \left(\frac{1}{9} \times \frac{9}{40}\right) + \left(\frac{1}{16} \times \frac{12}{40}\right) + \left(\frac{1}{25} \times \frac{5}{20}\right)$ $= \frac{133}{800} \quad (0.16625)$ $\text{Var}\left(\frac{1}{X}\right) = \frac{133}{800} - \frac{49}{400}$ $= \frac{7}{160}$ | M1 A1 m1 Adep1 | 4 | At least 4 of these terms added (accept decimal equivalents) (can be implied by $\frac{133}{800}$ seen with no other working shown) $\left[\text{their } E\left(\frac{1}{X^2}\right) \right] - \left(\frac{7}{20}\right)^2$ AG (allow 0.04375 seen) | | | | | | | | | | | | | | | | | | |
| (c)(i) | <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>y</td> <td>40</td> <td>20</td> <td>13$\frac{1}{3}$</td> <td>10</td> <td>8</td> </tr> <tr> <td>p</td> <td>$\frac{3}{40}$</td> <td>$\frac{6}{40}$</td> <td>$\frac{9}{40}$</td> <td>$\frac{12}{40}$</td> <td>$\frac{10}{40}$</td> </tr> </table> <p>Identifying $X = (2), 3, 4, 5$ or $Y = (20), 13\frac{1}{3}, 10, 8$</p> $P(X > 2) = \frac{9}{40} + \frac{12}{40} + \frac{5}{20}$ $= P(Y < 20)$ $= \frac{31}{40} \quad (0.775)$ | x | 1 | 2 | 3 | 4 | 5 | y | 40 | 20 | 13 $\frac{1}{3}$ | 10 | 8 | p | $\frac{3}{40}$ | $\frac{6}{40}$ | $\frac{9}{40}$ | $\frac{12}{40}$ | $\frac{10}{40}$ | M1 A1 A1 | 3 | Alternative: $Y < 20 \Rightarrow \frac{40}{X} < 20 \Rightarrow 40 < 20X \Rightarrow X > 2$ M1 (allow $<$ or \leq and $>$ or \geq in above) $P(Y < 20) = P(X > 2)$ $= 1 - \left(\frac{3}{40} + \frac{6}{40}\right) \quad \text{A1}$ $= \frac{31}{40} \quad (0.775) \quad \text{A1}$ |
| x | 1 | 2 | 3 | 4 | 5 | | | | | | | | | | | | | | | | | |
| y | 40 | 20 | 13 $\frac{1}{3}$ | 10 | 8 | | | | | | | | | | | | | | | | | |
| p | $\frac{3}{40}$ | $\frac{6}{40}$ | $\frac{9}{40}$ | $\frac{12}{40}$ | $\frac{10}{40}$ | | | | | | | | | | | | | | | | | |
| (ii) | $\frac{9}{40}$ seen irrespective of labelling $P(X < 4 Y < 20) = \frac{\frac{9}{40}}{\frac{31}{40}} = \frac{0.225}{0.775}$ $= \frac{9}{31} \quad (0.290)$ | B1 M1 A1 | 3 | As numerator or final answer (0.225) $= \frac{9}{40}$ (their (c)(i)) (or correct use of table) AFWF 0.29 to 0.2904 | | | | | | | | | | | | | | | | | | |
| Total | | | 14 | | | | | | | | | | | | | | | | | | | |

MS2B (cont)

| Q | Solution | Marks | Total | Comments | |
|-------|---|--|-----------|---|-------------------------------|
| 5(a) | $Y \sim N(\mu_y, 640^2)$ $n = 25$ and $\bar{y} = 19700$ $H_0: \mu_y = 20000$ $H_1: \mu_y \neq 20000$ (both) | B1 | | Alternative: $P(\bar{Y} < 19700) = P(Z < -2.34375)$ $= 1 - 0.99036$ $= 0.00964 \geq 0.005$ Accept H_0 | |
| | $\bar{Y} \sim N\left(20000, \frac{640^2}{25}\right)$ $z = \frac{19700 - 20000}{640/\sqrt{25}}$ $= -2.34375$ | M1 | | (-2.35 to -2.34) | |
| | $z_{crit} = \pm 2.5758$ | A1 | | (±2.57 to ±2.58) | |
| | Accept H_0 | B1 | | Use of $t \Rightarrow$ max B1M1A1 | |
| | Insufficient / no evidence (to suggest) that the mean (lifetime) has changed (from 20000 hours) | Adep1 | | dep on B1M1B1 | |
| | Mean (lifetime) has not changed at 1% level (of significance) | Edep1 | 6 | dep on Adep1 | |
| | | | | If incorrect hypotheses then B0 \Rightarrow max M1A1B1 ie final Adep1Edep1 not available | |
| | (b)(i) | $\mu < 10000$ | B1 | 1 | |
| | (ii) | $n = 16$ and $s = 500$; $t_{crit} = 1.753$ $sd(\bar{X}) = \frac{500}{\sqrt{16}}$ (125) Critical value is one of: $10000 \pm 1.753 \times \frac{500}{\sqrt{16}}$ (considered) | B1 | | For t_{crit} (ignore signs) |
| | | Choose 9780 (3sf) | B1 | | Ignore notation |
| | (\Rightarrow critical region: $\bar{x} < 9780$) | M1 | | M0 if only considered upper value No ft on incorrect t value | |
| | \therefore Range of values for \bar{x} which leads Christine not to reject $H_0: \mu = 10000$ is: $\bar{x} > 9780$ | A1 | 5 | AWFW 9780 to 9781 (ignore inequality) If z used then max B0B1M0A0A0 | |
| (iii) | No error | A1 | 1 | Allow $\bar{x} \geq 9780$ to 9781 | |
| | Total | | 13 | Ignore any subsequent statements | |

MS2B (cont)

| Q | Solution | Marks | Total | Comments |
|--|--|-------|--|---|
| 6(a) | $F(x) = \int \frac{3}{8}(x^2 + 1) dx$ | M1 | | Ignore limits |
| | $= \frac{3}{8} \left[\frac{x^3}{3} + x \right]$ or $= \frac{1}{8}x^3 + \frac{3}{8}x$ | A1 | | Either |
| | $= \frac{1}{8}x(x^2 + 3)$ | A1 | 3 | (including use of correct limits 0 and x or $+c$ used and evaluated) (AG) |
| (b) | $F(m) = \frac{1}{2}$ | B1 | | |
| | $F(1) = \frac{1}{8} \times 1 \times 4 = \frac{1}{2}$ | B1 | 2 | AG |
| (c) | Upper quartile lies in range $1 < x < 2$ such that $F(q) = \frac{3}{4}$ | | | $\frac{1}{2} + \int_1^q \frac{1}{4}(5 - 2x) dx = \frac{3}{4}$ |
| | $\int_1^q \frac{1}{4}(5 - 2x) dx = \frac{1}{4}$ | M1 | | Alternative: $\int_q^2 \frac{1}{4}(5 - 2x) dx = \frac{1}{4}$ |
| | $[5x - x^2]_1^q = 1$ | | | $[5x - x^2]_q^2 = 1$ |
| | $5q - q^2 - 4 = 1$ | | | $(10 - 4) - (5q - q^2) = 1$ |
| | $q^2 - 5q + 5 = 0$ | A1 | | $6 - 5q + q^2 = 1$ $q^2 - 5q + 5 = 0$ |
| | $q = \frac{5 \pm \sqrt{25 - 20}}{2}$ or $\frac{1}{2}(5 \pm \sqrt{5})$ | M1 | | Correct use of formula (OE) to give the two surd answers to given quadratic equation |
| | but $1 < q < 2$ [or ($q < 2$)] | m1 | | |
| $\therefore q = \frac{1}{2}(5 - \sqrt{5})$ | A1 | 5 | Must qualify with a numerical comparison, not just quote the given answer; dep on M1; AG | |
| (d) | $P(X > 1.5) = \frac{1}{2} \left[\frac{1}{2} + \frac{1}{4} \right] \times \frac{1}{2}$ | M1 | | $P(X < 1.5) = 0.5 + \frac{1}{2} \left[\frac{3}{4} + \frac{1}{2} \right] \times \frac{1}{2}$ (M1) |
| | $= \frac{3}{16}$ (0.1875) | A1 | | $= \frac{1}{2} + \frac{1}{2} \times \frac{5}{4} \times \frac{1}{2}$ |
| | $P(X > q) = \frac{1}{4}$ (0.25) | B1 | | $= \frac{1}{2} + \frac{5}{16} = \frac{13}{16}$ (A1) |
| | $P(q < X < 1.5) = \frac{1}{4} - \frac{3}{16}$ | | | $P(X < q) = \frac{3}{4}$ (0.75) (B1) |
| $= \frac{1}{16}$ (0.0625) | A1 | 4 | $P(q < X < 1.5) = \frac{13}{16} - \frac{3}{4} = \frac{1}{16}$ (A1) (0.0625) | |

MS2B (cont)

| Q | Solution | Marks | Total | Comments |
|------------------|---|-------|-----------|---|
| <p>6(d) cont</p> | <p>OR</p> $\int_{1.5}^2 \frac{1}{4}(5-2x) dx = \frac{3}{16} \text{ etc (M1A1)}$ <p>NB statement $F(1.5) - \frac{3}{4} = \frac{1}{16}$ (OE) scores 4 marks</p> <p>Alternative:</p> $\int_q^{1.5} \frac{1}{4}(5-2x) dx = \left[-\frac{1}{16}(5-2x)^2 \right]_{\frac{5-\sqrt{5}}{2}}^{1.5}$ <p style="text-align: right;">(M1)</p> $= -\frac{1}{16}(4) - \left[-\frac{1}{16}(\sqrt{5})^2 \right] \text{ (sub) (A1)}$ $= -\frac{4}{16} + \frac{5}{16} \text{ (A1)}$ $= \frac{1}{16} \text{ (A1)}$ | | | <p>OR</p> $\int_q^{1.5} \frac{1}{4}(5-2x) dx = \frac{1}{4} [5x - x^2]_q^{1.5} \text{ (M1)}$ <p>(correct integration and limits) Allow use of $q = 1.38$ to $q = 1.382$ in limits for M1 Whatever follows must be exact</p> $= \frac{1}{4} [(7.5 - 2.25) - (5q - q^2)] \text{ (A1)}$ <p>for use of $5q - q^2 = 5$ or showing $5q - q^2 = 5$ by substituting $q = \frac{1}{2}(5 - \sqrt{5})$ (A1)</p> $= \frac{1}{4} [5.25 - 5] = \frac{1}{16} \text{ (A1)}$ <p>Alternative using F(x):</p> <p>for $1 \leq x \leq 2$</p> $F(x) = \frac{1}{2} + \int_1^x \frac{1}{4}(5-2x) dx$ $= \frac{1}{2} + \frac{1}{4} [5x - x^2]_1^x$ $= \frac{1}{2} + \frac{1}{4} [(5x - x^2) - (5 - 1)]$ $= \frac{1}{4} (2 + 5x - x^2 - 4)$ $= \frac{1}{4} (5x - x^2 - 2) \text{ (seen or used) (M1)}$ $F(1.5) = \frac{1}{4} (7.5 - 2.25 - 2) = \frac{3.25}{4}$ $= 0.8125 = \frac{13}{16} \text{ (A1)}$ $F(q) = \frac{1}{16} (50 - 10\sqrt{5} - (25 - 10\sqrt{5} + 5) - 8)$ $= \frac{12}{16} \text{ OE (B1)}$ $P(q < X < 1.5) = \frac{13}{16} - \frac{12}{16} = \frac{1}{16} \text{ (A1)}$ |
| | Total | | 14 | |
| | TOTAL | | 75 | |