



**General Certificate of Education (A-level)  
January 2012**

**Mathematics**

**MFP3**

**(Specification 6360)**

**Further Pure 3**

**Final**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2012 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

Q	Solution	Marks	Total	Comments
1(a)	$y(1.1) = y(1) + 0.1 \left[ \frac{2-1}{4+1} \right]$	M1A1	3	
	$= 2 + 0.02 = 2.02$	A1		
(b)	$y(1.2) = y(1) + 2(0.1)\{f[1.1, y(1.1)]\}$	M1	3	ft on c's answer to (a) CAO Must be 2.036
	$= 2 + 2(0.1) \left[ \frac{2.02-1.1}{2.02^2+1.1} \right]$	A1F		
	$= 2.035518... = 2.036$ to 3dp	A1		
<b>Total</b>			<b>6</b>	
2	$\sqrt{4+x} = 2 \left( 1 + \frac{x}{4} \right)^{\frac{1}{2}} = 2 \left[ 1 + \frac{1}{2} \left( \frac{x}{4} \right) + O(x^2) \right]$	M1	3	Attempt to use binomial theorem OE The notation $O(x^n)$ can be replaced by a term of the form $kx^n$  Division by $x$ stage before taking the limit  CSO NMS 0/3
	$\left[ \frac{\sqrt{4+x}-2}{x+x^2} \right] = \left[ \frac{\frac{x}{4} + O(x^2)}{x+x^2} \right] = \left[ \frac{\frac{1}{4} + O(x)}{1+x} \right]$	m1		
	$\lim_{x \rightarrow 0} \left[ \frac{\sqrt{4+x}-2}{x+x^2} \right] = \frac{1}{4}$	A1		
<b>Total</b>			<b>3</b>	
3	$m^2 + 2m + 10 = 0$	M1	10	PI  OE Ft on incorrect <b>complex value</b> of $m$  c's CF+ c's non-zero PI but must have 2 arb consts  ft c's $k$ ie $A = 5 - k, k \neq 0$  Attempt to differentiate c's <b>GS</b> (ie CF + PI)  CSO
	$m = -1 \pm 3i$	A1		
	Complementary function is $(y \Rightarrow) e^{-x} (A \cos 3x + B \sin 3x)$	A1F		
	Particular integral: try $y = ke^x$ $k + 2k + 10k = 26 \Rightarrow k = 2$	M1 A1		
	(GS $y \Rightarrow) e^{-x} (A \cos 3x + B \sin 3x) + 2e^x$	B1F		
	$x = 0, y = 5 \Rightarrow 5 = A + 2$ so $A = 3$	B1F		
	$\frac{dy}{dx} =$ $e^{-x}(-3A \sin 3x + 3B \cos 3x - A \cos 3x - B \sin 3x) + 2e^x$	M1		
$11 = 3B - A + 2$ ( $B = 4$ )	A1			
$y = e^{-x} (3 \cos 3x + 4 \sin 3x) + 2e^x$	A1			
<b>Total</b>			<b>10</b>	

Q	Solution	Marks	Total	Comments
4(a)	IF is $\exp\left(\int \frac{2}{x} dx\right)$	M1	7	and with integration attempted
	$= e^{2\ln x}$	A1		PI
	$= x^2$	A1		
	$\frac{d}{dx}[yx^2] = x^2 \ln x$	M1		LHS; PI
	$\Rightarrow yx^2 = \int (\ln x) \frac{d}{dx}\left(\frac{x^3}{3}\right)$	M1		Attempt integration by parts in correct direction to integrate $x^p \ln x$
	$= \frac{x^3}{3} \ln x - \int \frac{x^2}{3} dx$	A1		RHS
(b)	Now, as $x \rightarrow 0$ , $x^k \ln x \rightarrow 0$	E1	3	Must be stated explicitly for a value of $k > 0$
	As $x \rightarrow 0$ , $y \rightarrow 0 \Rightarrow A = 0$	B1		Const of int = 0 must be convincing
	$yx^2 = \frac{x^3}{3} \ln x - \frac{x^3}{9}$			
	When $x = 1$ , $y = -\frac{1}{9}$	B1F		ft on one slip but must have made a realistic attempt to find A
	<b>Total</b>		<b>10</b>	

Q	Solution	Marks	Total	Comments
5(a)	The interval of integration is infinite	E1	1	OE
(b)	$u = x^2 e^{-4x} + 3 \Rightarrow du = (2xe^{-4x} - 4x^2 e^{-4x}) dx$ $\int \frac{x(1-2x)}{x^2 + 3e^{4x}} dx = \int \frac{1}{2} \times \frac{2x(1-2x)e^{-4x}}{x^2 e^{-4x} + 3} dx$ $= \frac{1}{2} \times \int \frac{1}{u} du$ $= \frac{1}{2} \ln u + c = \frac{1}{2} \ln(x^2 e^{-4x} + 3) \{+c\}$	M1  A1 A1	3	du/dx or 'better'  OE Condone missing $c$ . Accept later substitution back if explicit
(c)	$I = \int_{\frac{1}{2}}^{\infty} \frac{x(1-2x)}{x^2 + 3e^{4x}} dx$ $= \lim_{a \rightarrow \infty} \int_{\frac{1}{2}}^a \frac{x(1-2x)}{x^2 + 3e^{4x}} dx$ $= \lim_{a \rightarrow \infty} \frac{1}{2} \left\{ \ln(a^2 e^{-4a} + 3) - \ln\left(\frac{e^{-2}}{4} + 3\right) \right\}$ $= \frac{1}{2} \ln \left\{ \lim_{a \rightarrow \infty} (a^2 e^{-4a} + 3) \right\} - \frac{1}{2} \ln\left(\frac{e^{-2}}{4} + 3\right)$ <p>Now <math>\lim_{a \rightarrow \infty} (a^2 e^{-4a}) = 0</math></p> $I = \frac{1}{2} \ln 3 - \frac{1}{2} \ln\left(\frac{e^{-2}}{4} + 3\right)$	M1  M1  E1 A1	4	Uses part (b) and $F(a) - F(1/2)$  Stated explicitly (could be in general form) CSO ACF
	<b>Total</b>		<b>8</b>	

Q	Solution	Marks	Total	Comments
6(a)	$y = \ln \cos 2x \Rightarrow y'(x) = \frac{1}{\cos 2x} (-2 \sin 2x)$	M1 A1	6	Chain rule
	$y''(x) = -4 \sec^2 2x$	m1		$\lambda \sec^2 2x$ OE
	$y'''(x) = -8 \sec 2x (2 \sec 2x \tan 2x)$	M1		$K \sec^2 2x \tan 2x$ OE
	$\{y'''(x) = -16 \tan 2x (\sec^2 2x)\}$			
	$y''''(x) = -16[2 \sec^2 2x (\sec^2 2x) + \tan 2x (2 \sec 2x (2 \sec 2x \tan 2x))]$	M1 A1		Product rule OE ACF
(b)	$y(0) = 0, y'(0) = 0, y''(0) = -4, y'''(0) = 0, y''''(0) = -32$	B1F		ft c's derivatives
	$\ln \cos 2x \approx 0 + 0 + \frac{x^2}{2}(-4) + 0 + \frac{x^4}{4!}(-32)$ $\approx -2x^2 - \frac{4}{3}x^4$	M1  A1	3	CSO throughout parts (a) and (b) AG
(c)	$\ln(\sec^2 2x) = -2 \ln(\cos 2x)$	M1		PI
	$\approx 4x^2 + \frac{8}{3}x^4$	A1	2	
<b>Total</b>			<b>11</b>	

Q	Solution	Marks	Total	Comments
7(a)	$u = xy$ $\frac{du}{dx} = y + x \frac{dy}{dx}$ $\frac{d^2u}{dx^2} = \frac{dy}{dx} + \left( \frac{dy}{dx} + x \frac{d^2y}{dx^2} \right)$ $x \frac{d^2y}{dx^2} + 2(3x+1) \frac{dy}{dx} + 3y(3x+2) = 18x$ $\left( x \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} \right) + 6 \left( x \frac{dy}{dx} + y \right) + 9xy = 18x$ $\frac{d^2u}{dx^2} + 6 \frac{du}{dx} + 9u = 18x$	M1 A1  A1		Product rule OE OE OE
		A1	4	CSO AG Be convinced
(b)	$\frac{d^2u}{dx^2} + 6 \frac{du}{dx} + 9u = 18x$ CF: Aux eqn $m^2 + 6m + 9 = 0$ $(m+3)^2 = 0$ so $m = -3$ CF: $(u =) e^{-3x} (Ax + B)$ PI: Try $(u =) px + q$ $0 + 6p + 9(px + q) = 18x$ $9p = 18, \quad 6p + 9q = 0$ $p = 2; \quad q = -\frac{12}{9}$ $u = e^{-3x} (Ax + B) + 2x - \frac{4}{3}$ $xy = e^{-3x} (Ax + B) + 2x - \frac{4}{3}$ $y = \frac{1}{x} \left\{ e^{-3x} (Ax + B) + 2x - \frac{4}{3} \right\}$	M1 A1 A1F  M1 m1 A1		PI PI  PI. Must be more than just stated  Both
		B1F		c's CF + c's PI but must have 2 constants, also must be in the form $u = f(x)$
		A1	8	
	<b>Total</b>		<b>12</b>	



Q	Solution	Marks	Total	Comments
8(a)	Area = $\frac{1}{2} \int (3 + 2 \cos \theta)^2 d\theta$	M1	6	Use of $\frac{1}{2} \int r^2 d\theta$ or $\int_0^\pi r^2 d\theta$
	$= \frac{1}{2} \int_0^{2\pi} (9 + 12 \cos \theta + 4 \cos^2 \theta) d\theta$ $= \int_0^{2\pi} (4.5 + 6 \cos \theta + (1 + \cos 2\theta)) d\theta$ $= \left[ 4.5\theta + 6 \sin \theta + \theta + \frac{1}{2} \sin 2\theta \right]_0^{2\pi}$ $= 11\pi$	B1 B1 M1 A1F A1		Correct expn of $[3 + 2 \cos \theta]^2$ Correct limits Attempt to write $\cos^2 \theta$ in terms of $\cos 2\theta$ Correct integration ft wrong coefficients CSO
(b)(i)	$x^2 + y^2 - 8x + 16 = 16$	M1	6	Use of <b>any two</b> of $x = r \cos \theta$ , $y = r \sin \theta$ , $x^2 + y^2 = r^2$
	$r^2 - 8r \cos \theta + 16 = 16 \Rightarrow r = 8 \cos \theta$	A1		
	At intersection, $8 \cos \theta = 3 + 2 \cos \theta$ $\Rightarrow \cos \theta = \frac{3}{6}$	M1		Equating $rs$ or equating $\cos \theta$ s with a further step to solve eqn. (OE eg $4r = 12 + r \Rightarrow 4r - r = 12$ )
	Points $\left(4, \frac{\pi}{3}\right)$ and $\left(4, \frac{5\pi}{3}\right)$	A1		OE
	$AB = 2 \times \left(4 \sin \frac{\pi}{3}\right)$ $= 4\sqrt{3}$	M1 A1		Valid method to find $AB$ , ft c's $r$ and $\theta$ values OE surd
(ii)	Let $M$ =centre of circle then $\angle AMB = \frac{2\pi}{3}$	B1	3	Accept equiv eg $\angle AMO = \frac{\pi}{3}$
Length of arc $AOB$ of circle = $4 \times \frac{2\pi}{3}$	M1	Use of arc = $4 \times (\angle AMB \text{ in rads})$		
Perimeter of segment $AOB = \frac{8\pi}{3} + 4\sqrt{3}$	A1			
	<b>Total</b>		<b>15</b>	
	<b>Alternative to (b)(i):</b> Writing $r = 3 + 2 \cos \theta$ in cartesian form (M1A1) Finding cartesian coordinates of points $A$ and $B$ ie $(2, \pm 2\sqrt{2})$ (M1A1) Finding length $AB$ (M1A1)			
	<b>TOTAL</b>		<b>75</b>	