



**General Certificate of Education (A-level)
June 2013**

Mathematics

MFP1

(Specification 6360)

Further Pure 1

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Marks	Total	Comments
1	$(x_2 =) 10 - \frac{(10^3 - 10^2 + 4 \times 10 - 900)}{(3 \times 10^2 - 2 \times 10 + 4)}$	B1		$10 - \frac{f(10)}{f'(10)}$ with a correct numerical expression or value PI for $f(10)$.
	$\left(= 10 - \frac{1000 - 100 + 40 - 900}{300 - 20 + 4} \right)$	B1		$10 - \frac{f(10)}{f'(10)}$ with a correct numerical expression or value PI for $f'(10)$.
	$\left(= 10 - \frac{40}{284} = 10 - 0.1408... \right)$ $(= 9.85915....) = 9.859$ (to 4 sf)	B1	3	Must be 9.859
	Total		3	
2(a)(i)	$\mathbf{A} - \mathbf{B} = \begin{bmatrix} p-3 & 1 \\ 2 & p-3 \end{bmatrix}$	B1	1	
(ii)	$\mathbf{AB} = \begin{bmatrix} p & 2 \\ 4 & p \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 3p+4 & p+6 \\ 12+2p & 4+3p \end{bmatrix}$	M1		Finding \mathbf{AB} and at least 2 elements correct
		A1	2	CSO
(b)	$\mathbf{A} - \mathbf{B} + \mathbf{AB} = \begin{bmatrix} 4p+1 & p+7 \\ 14+2p & 1+4p \end{bmatrix}$	B1F		Only ft if all matrices are 2 by 2 PI by later correct work
	$\mathbf{A} - \mathbf{B} + \mathbf{AB} = k \mathbf{I} = k \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	B1		\mathbf{I} used as or equated to $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ at some stage
	$(p+7=0, 14+2p=0 \Rightarrow) p=-7$	B1		$p=-7$ provided it gives the relevant two zero elements
	$p=-7 \Rightarrow \mathbf{A} - \mathbf{B} + \mathbf{AB} = \begin{bmatrix} -27 & 0 \\ 0 & -27 \end{bmatrix} = -27 \mathbf{I}$ $\Rightarrow k = -27$	B1	4	CSO Either $-27\mathbf{I}$ (no earlier errors) for B1 OR $k = -27$ with either $\begin{bmatrix} -27 & 0 \\ 0 & -27 \end{bmatrix}$ or $27 \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$ or $-27 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ seen before (no earlier errors) for B1
	Total		7	

Q	Solution	Marks	Total	Comments
3(a)	$\cos(5x + 40^\circ) = \cos 65^\circ$ $5x + 40^\circ = \pm 65^\circ$ $5x + 40^\circ = 360n^\circ + 65^\circ, 5x + 40^\circ = 360n^\circ - 65^\circ$ $x = \frac{360n^\circ + 65^\circ - 40^\circ}{5}, x = \frac{360n^\circ - 65^\circ - 40^\circ}{5}$ $x = 72n^\circ + 5^\circ, x = 72n^\circ - 21^\circ$	<p>B1</p> <p>M1</p> <p>m1</p> <p>A2,1,0</p>	5	<p>Both $\pm 65^\circ$ OE eg $5x + 40 = 65, 295$</p> <p>$5x + 40 = 360n \pm \alpha$ Either one, OE Condone $2n\pi$ for $360n$</p> <p>Either one, OE Correct rearrangement of $5x + 40 = 360n \pm \alpha$ OE to $x =$. Condone $2n\pi$ for $360n$</p> <p>OE Full set of correct solns. in degrees written in a simplified form. (A1 if not in a simplified form) (A0 if radians present in answer)</p>
(b)	$\frac{\sqrt{3}-1}{2\sqrt{2}} = (\cos \frac{\pi}{4})[\cos(a\pi) + \cos(b\pi)]$ $\frac{1}{\sqrt{2}} \left(\frac{\sqrt{3}}{2} - \frac{1}{2} \right) = \frac{1}{\sqrt{2}} [\cos(a\pi) + \cos(b\pi)]$ $\cos a\pi = \frac{\sqrt{3}}{2} = \cos \frac{\pi}{6}, (a = \frac{1}{6})$ $\cos b\pi = -\frac{1}{2} = \cos \frac{2\pi}{3}, (b = \frac{2}{3})$ $\sin \frac{\pi}{12} = \cos \left(\frac{\pi}{4} \right) \left[\cos \left(\frac{1}{6} \pi \right) + \cos \left(\frac{2}{3} \pi \right) \right]$	<p>B1</p> <p>B1</p> <p>B1</p>	3	<p>Recognising $\cos \frac{\pi}{4} = \frac{\sqrt{2}}{2}$ (or $= \frac{1}{\sqrt{2}}$)</p> <p>PI eg by seeing $\frac{1}{\sqrt{2}} \left(\frac{\sqrt{3}}{2} - \frac{1}{2} \right)$</p> <p>OE ie any correct rational value for a which satisfies $\cos a\pi = \frac{\sqrt{3}}{2}$</p> <p>OE ie any correct rational value for b which satisfies $\cos b\pi = -\frac{1}{2}$</p> <p>Note: labels a and b could be interchanged.</p>
	Total		8	

Q	Solution	Marks	Total	Comments
4(a)(i)	$(z - 2i)^* = (x + yi - 2i)^* = x + (2 - y)i$	B1	1	$x + 2i - yi$ OE rearrangement
(ii)	$(z - 2i)^* = 4i z + 3 = 4ix + 4i^2y + 3 = 4ix - 4y + 3$ $x + (2 - y)i = 4ix - 4y + 3$ (#) Real parts: $x = -4y + 3$ Imaginary parts: $2 - y = 4x$	B1		$i^2 = -1$ used
		M1		Attempting to equate, without mixing real and imaginary terms, both the real parts and the imag. parts for the c's eqn (#).
		A1F		If not corrected, ft on [c's(a)(i)] = $4ix - 4y + 3$ provided both the resulting linear equations have non zero x, y and const terms
	$y = \frac{2}{3}, x = \frac{1}{3}$	A1		Solving correct equations, to obtain either $x = \frac{1}{3}$ OE or $y = \frac{2}{3}$ OE
	$(z =) \frac{1}{3} + \frac{2}{3}i$	A1	5	$\frac{1}{3} + \frac{2}{3}i$
(b)	(One of the) coefficients (of the quadratic equation is) not real.	E1	1	OE eg Sum of roots is $-10i$ so p cannot be real if roots are $p \pm qi$
	Total		7	

Q	Solution	Marks	Total	Comments
5(a)(i)	$y = 2x^2 - 5x$ $y_Q = 2(1+h)^2 - 5(1+h) = 2 + 4h + 2h^2 - 5 - 5h$ $(= 2h^2 - h - 3)$	B1		$y_Q = 2(1+h)^2 - 5(1+h)$ with correct expansion of brackets PI.
	$\text{Grad.} = \frac{y_Q - y_P}{x_Q - x_P} = \frac{2(1+h)^2 - 5(1+h) - (-3)}{1+h-1}$	M1		Use of correct formula for gradient
	$= \frac{2h^2 - h - 3 - (-3)}{h} = \frac{2h^2 - h}{h} = 2h - 1$	A1	3	CSO
(ii)	As $h \rightarrow 0$, (grad of $PQ \rightarrow$ grad of tangent at P)	E1		$h = 0$ scores E0
	(ie) gradient (of tangent at P) = -1 Now gradient of $x+y=0$ (or $y=-x$) is also -1 \Rightarrow tangent at P is parallel to line $x+y=0$	E1	2	Dep on $h \rightarrow 0$ or $h=0$ being used earlier
(b)	$I = \int_1^{\infty} x^{-4}(2x^2 - 5x) dx = \int_1^{\infty} (2x^{-2} - 5x^{-3}) dx$			
	$I = \left[-2x^{-1} - 5 \frac{x^{-2}}{-2} \right]_1^{\infty}$	M1		At least one term correct
	As $x \rightarrow \infty$, $x^{-1} \rightarrow 0$ and $x^{-2} \rightarrow 0$	E1		OE Ft on kx^{-n} provided M1 awarded
	$I = 0 - (-2 + 5/2) = -\frac{1}{2}$	A1	3	(I =) $-\frac{1}{2}$ Dep on both terms integrated correctly in the M1 line
	Total		8	

Q	Solution	Marks	Total	Comments	
6(a)	$\alpha + \beta = -\frac{3}{2}$	B1	2	OE	
	$\alpha\beta = -3$	B1		OE	
	(b)	$\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$	M1	3	Using correct identity for $\alpha^3 + \beta^3$ in terms of $\alpha + \beta$ and $\alpha\beta$.
		$= \left(-\frac{3}{2}\right)^3 - 3(-3)\left(-\frac{3}{2}\right)$	A1F		with ft/or correct substitution
		$= -\frac{27}{8} - \frac{27}{2} = -\frac{135}{8}$	A1		CSO AG. Correct evaluation of each of $(-1.5)^3$ and $-3(-3)(-1.5)$ must be seen before the printed answer is stated
	(c)	Sum = $\alpha + \frac{\alpha}{\beta^2} + \beta + \frac{\beta}{\alpha^2}$	M1	6	Writing $\alpha + \frac{\alpha}{\beta^2} + \beta + \frac{\beta}{\alpha^2}$ in a suitable form with ft/or correct substitution
		$= \alpha + \beta + \frac{\alpha^3 + \beta^3}{(\alpha\beta)^2} = -\frac{3}{2} + \frac{-135/8}{9}$			
		Sum = $-\frac{27}{8}$	A1		PI OE exact value eg -3.375 (A0 if $\alpha\beta = 3$ used to get $(\alpha\beta)^2 = 9$)
		Product = $\alpha\beta + \frac{\beta}{\alpha} + \frac{\alpha}{\beta} + \frac{1}{\alpha\beta}$	M1		PI OE exact value eg -3.375 (A0 if $\alpha\beta = 3$ used to get $(\alpha\beta)^2 = 9$)
		$= \alpha\beta + \frac{\alpha^2 + \beta^2}{\alpha\beta} + \frac{1}{\alpha\beta}$ (*)			
Now $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$ $(= \frac{9}{4} + 6)$					
Product = $-3 - \frac{1}{3}\left(\frac{9}{4} + 6\right) - \frac{1}{3} = -\frac{73}{12}$	A1	PI OE exact value			
$x^2 - Sx + P (= 0)$	M1	Using correct general form of LHS of eqn with ft substitution of c's S and P values.			
Eqn is $24x^2 + 81x - 146 = 0$	A1	OE but integer coefficients and '=' needed			
Total			11		

Q	Solution	Marks	Total	Comments
7(a)	$f(x) = 4x^3 - x - 540\,000$ $f(51) = -9447 (<0)$; $f(52) = 22380 (>0)$; Since sign change (and f continuous), $51 < \alpha < 52$	M1 A1	2	$f(51)$ and $f(52)$ both considered All values and working correct plus relevant concluding statement involving '51' and '52'.
(b)(i)	$S_n = \sum_{r=1}^n (2r-1)^2 = \sum 4r^2 - \sum 4r + \sum 1$ $= 4 \frac{n}{6} (n+1)(2n+1) - 4 \frac{n}{2} (n+1) + \sum_{r=1}^n 1$ $= 4 \frac{n}{6} (n+1)(2n+1) - 4 \frac{n}{2} (n+1) + n$ $= \frac{n}{3} [2(2n^2 + 3n + 1) - 6(n+1) + 3] = \frac{n}{3} [4n^2 - 1]$	M1 m1 B1 A1	5	Splitting up the sum into separate sums. PI by m1 line below or better Substitution of correct formulae from FB for the two summations B1 for $\sum_{r=1}^n 1 = n$ stated or used CSO
(ii)	$(6 S_n = 2n[4n^2 - 1]) = 2n(2n-1)(2n+1)$ $(2n-1)$, $2n$ and $(2n+1)$ are consecutive integers	B1 E1	2	Terms in any order Terms must be identified and statement 'consecutive integers'
(c)	$S_n = 1^2 + 3^2 + 5^2 + \dots + (2n-1)^2$ ie sum of squares of first n odd numbers so need least N such that $S_N > 180\,000$ $S_{52} = \frac{52}{3} [4 \times 52^2 - 1] = 187460$ and $S_{51} = 176851$ Smallest value of N is 52	M1 A1	2	Either $\frac{n}{3} [4n^2 - 1] = 180000$ or $2N(2N-1)(2N+1) = 1080000$ or S_{52} and S_{51} both attempted (or = replaced by > or by \geq) CSO Fully and correctly justified. NMS $N=52$ scores 0/2
Total			11	

Q	Solution	Marks	Total	Comments
8(a)	$\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$	M1		Matrix in form $\begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}$, where $\lambda \neq 0, \mu \neq 0$ and $\lambda \neq \mu$
(b)(i)	$y = \sqrt{3}x = \tan 60^\circ x \quad \begin{bmatrix} \cos 120^\circ & \sin 120^\circ \\ \sin 120^\circ & -\cos 120^\circ \end{bmatrix}$ Required matrix is $\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$	M1	2	$\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ $\begin{bmatrix} \cos 120^\circ & \sin 120^\circ \\ \sin 120^\circ & -\cos 120^\circ \end{bmatrix}$ PI For M mark, condone dec approx 0.86 or 0.87 or better in place of $\sin 120^\circ$
(ii)	$\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix} = \dots\dots$ $= \begin{bmatrix} \frac{1}{2} & \frac{3\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{3}{2} \end{bmatrix}$	M1	2	Attempt to multiply c's (b)(i) 2by2 matrix and c's (a) 2by2 matrix in correct order. OE but must be in exact surd form.
	Total		6	

Q	Solution	Marks	Total	Comments
9(a)	(HA) $y = 1$ (VA) $x^2 - 2x - 3 = 0$ $(x - 3)(x + 1) = 0$ $x = -1$ and $x = 3$	B1 M1 A1	3	$y = 1$ OE eqn PI OE eg use of quadratic formula Both needed OE eqn(s)
(b)(i)	$k = \frac{x^2 - 2x + 1}{x^2 - 2x - 3} \Rightarrow kx^2 - 2kx - 3k = x^2 - 2x + 1$ $kx^2 - 2kx - 3k - x^2 + 2x - 1 = 0$ $(k - 1)x^2 - 2(k - 1)x - (3k + 1) = 0$	B1	1	AG Must see the two stages, correct elimination of fraction and a correct rearrangement to $\dots = 0$, along with correct elimination of brackets before printed answer is stated.
(ii)	Discriminant $b^2 - 4ac$ $\{ 4(k - 1)^2 + 4(k - 1)(1 + 3k) \}$ Line intersects curve $\Rightarrow b^2 - 4ac \geq 0$ $\Rightarrow 4(k - 1)^2 + 4(k - 1)(1 + 3k) \geq 0$ $\Rightarrow 4(k - 1)[k - 1 + 1 + 3k] \geq 0$, $16k(k - 1) \geq 0$ ie $k^2 - k \geq 0$	M1 A1 A1	3	$b^2 - 4ac$, OE, in terms of k ; condoning one minor error in substitution. A correct inequality where k is the only unknown CSO AG Must be convinced
(iii)	$k^2 - k \geq 0$, $k(k - 1) \geq 0$, $k \leq 0$, $k \geq 1$ Critical values $k = 0$, $(k = 1)$ $k \neq 1$ since there is no point on the curve where $y = 1$ ($x^2 - 2x - 3 \neq x^2 - 2x + 1$) $k = 0$, $-x^2 + 2x - 1 = 0$ or $y = 0$, $x^2 - 2x + 1 = 0$ (Only one) stationary point (and its coordinates are) (1, 0)	B1 E1 M1 A1	4	For $k = 0$ either as an equation or inequality. OE Valid explanation, with no accuracy errors, to discount $k = 1$ OE 'stationary' with either (1, 0) or $\{x = 1, y = 0\}$
(c)		B1 B1 B1	3	Curve with three distinct branches Branch between VAs, correct shape, no part of the branch above the x -axis, only intersection with y -axis at a point below the origin, and its max pt on the positive x -axis Fully correct curve drawn with each branch correctly approaching its relevant asymptotes
	Total		14	
	TOTAL		75	