



AS

Mathematics

MS1B Statistics 1B

Final Mark Scheme

6360
June 2017

Version/Stage: v1.0

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

General Notes for MS1B

- GN1** There is no allowance for misreads (MR) or miscopies (MC) unless specifically stated in a question.
- GN2** In general, a correct answer (to accuracy required) without working scores full marks but an incorrect answer (or an answer not to required accuracy) scores no marks.
- GN3** In general, a correct answer (to accuracy required) without units scores full marks.
- GN4** When applying AFWF, a slightly inaccurate numerical answer that is subsequently rounded to fall within the accepted range cannot be awarded full marks.
- GN5** Where percentage equivalent answers are permitted in a question, then penalise by **one accuracy mark** at the first **correct** answer but only if no indication of percentage (eg %) is shown.
- GN6** In questions involving probabilities, do **not** award **accuracy** marks for answers given in the form of a ratio or odds such as 13/47 given as 13:47 or 13:34.
- GN7** Accept decimal answers, providing that they have **at least two** leading zeros, in the form $c \times 10^n$ (eg 0.00321 as 3.21×10^{-3}).
- GN8** **Where a candidate's response to a part of a question is simply to label the part (eg (d)(i)) with nothing else (ie no attempt at a solution), then this is still treated as a response and marked as 0 rather than NR. Also, deleted work, if not replaced, should be marked and not treated as NR.**

Specific Notes for MS1B

1. **Question 1**
In part(a), the equation is NOT required but, if simply quoted, can score marks.
2. **Question 2**
In part (b), for example, “LCL < 25 < UCL so agree (with claim)” scores 2 marks.
3. **Question 3**
In parts (b) & (c), note the ranges if using Bdep1, Bdep1 and Mdep1.
4. **Question 4**
In part (c), equivalent contextual explanations to “positive” and “negative” are NOT acceptable.
5. **Question 5**
In part (b)(i), be aware of Note 3 in part (b)(i) when then marking part (b)(ii)(B).
6. **Question 6**
In part (b)(i), the Special Cases are by no means rare events!
7. **Question 7**
In part (a)(iii), note the stricter conditions for the awarding of the method marks.
In part(b)(ii), there is NO follow-through from part(b)(i).

Q	Solution	Marks	Total	Comments
2 (a)	Sample is random	B1		Must include both emboldened words
	Durations are normal (ly distributed)	B1		Must include both emboldened words
Notes	1 "Sample (data/values/durations/(telephone) calls) is random and is normally distributed" ⇒ B1 B0 2 "Durations (of (telephone) calls) are random and normally distributed" ⇒ B0 B1 3 "Length of calls are independent and normal(ly distributed)" ⇒ B0 B1			
	95% (0.95) ⇒ $z = \underline{1.96}$ Mean / $\bar{x} = \underline{21.5}$ CI for μ is: $(20.5 \text{ to } 22.5 \text{ AFWW}) \pm \begin{pmatrix} 1.96 \text{ AWR T} \\ \text{or} \\ 1.64 \text{ to } 1.65 \text{ AFWW} \end{pmatrix} \times \frac{7.5}{\sqrt{10}}$ Hence $\underline{21.5 \pm (4.6 \text{ to } 4.7)}$ or $\underline{(16.8 \text{ to } 16.9, 26.1 \text{ to } 26.2)}$	B1 B1 M2,1 (-1 ee) Adep1	7	AWR T (1.95996) CAO (Ignore notation) M0 if CI is not an interval or is not of the form $\bar{x} \pm z \times \frac{7.5}{\sqrt{10}}$ CAO/AFWW (21.5 ± 4.64855) Dependent on award of M2 AFWW (16.85, 26.15)
Notes	1 If award of M0 is followed by a numerically correct CI ⇒ 2 solutions 2 Use of z-value of 1.28 to 1.29 or 2.05 to 2.06, 2.32 to 2.33 or 2.57 to 2.58 ⇒ 1 error but any other z/t values ⇒ M0 3 Use of $s^2(n-1) = 96.11$, $s(n-1) = 9.98$, $s^2(n) = 89.65$ or $s(n) = 9.47$ ⇒ M0			
(b)	Clear correct comparison of 25 with CI in (a) (eg 25 is within CI or interval) Agree with or accept claim or Claim is (likely to be) true/correct/right/valid/accurate/supported/reasonable/possible	BF1 Bdep1	2	Statement must include reference to 25 F on CI providing it includes 25 Must have found an interval in (a) but quoting values for CI or CLs is not required Ignore additional statements such as those in Notes 3 & 4 below \ OE; dependent on BF1 Do NOT ignore conflicting conclusions such as that in Note 5 below
	Notes	1 Statement must clearly indicate that "25 is within the interval" OE 2 Statements of the form "25 is below the UCL" or "25 is above the LCL" ⇒ BF0 3 Statements of the form "It/this/mean/value/duration/minutes/(telephone)calls/etc is within the interval" ⇒ BF0 4 Statements of the form "25 is within 95% of the data/values/durations/minutes/(telephone)calls/etc" ⇒ BF0 5 Statements of the form "25 is within the interval but it is towards the UCL so may/may not be true" (OE) ⇒ BF1 Bdep0		
		Total	9	

Q	Solution	Marks	Total	Comments
3 (a)	$\text{Mean} = \underline{7.35}$ $= \underline{7.3 \text{ to } 7.4}$ $\text{Var}(n) = \underline{10.5} \quad \text{or} \quad \text{Var}(n-1) = \underline{10.6}$ $\text{Var}(n \text{ or } n-1) = \underline{10.4 \text{ to } 10.7}$	B2 (B1) B2 (B1)	 4	CAO $(\sum x = 588)$ AFWW AWRW $(10.47750 \text{ or } 10.61013)$ $(\sum x^2 = 5160)$ AFWW
Notes	1 Mean = 182 to 182.4 \Rightarrow B0 B2 max 2 Answer quoted as 3.22 ² to 3.26 ² alone \Rightarrow B2 B1 max 3 Answer quoted as 3.22 to 3.26 alone \Rightarrow B2 B0 max 4 If answers not identified, then mark as mean followed by variance			
SC	1 If and only if B0 B0, then award M1 for at least 4 of 1, 3, 5, 7, 9, 11, 13 seen or for $\frac{(508 \text{ to } 668)}{80}$ or (6.35 to 8.35)			
(b)	$\text{Mean} = \underline{182 \text{ to } 182.4}$ <p>or</p> $\text{Mean} = \underline{175 + (\text{mean in (a)})}$ $\text{Var}(n \text{ or } n-1) = \underline{10.4 \text{ to } 10.7}$ <p>or</p> $\text{Var}(n \text{ or } n-1) = \underline{\text{value of Var stated in (a)}}$	B1 (Bdep1) B1 (Bdep1)	 2	AFWW; irrespective of value quoted/stated as mean in (a) Evaluated (at least 3sf) using value quoted/stated as mean in (a) and dep on 6 < Mean < 9 in (a) AFWW; irrespective of value quoted/stated as variance in (a) Must be identical (at least 3 sf) to value quoted/stated as variance in (a) and dep on 9 < Var < 12 in (a)
(c)	$\text{Mean} = \underline{5.97 \text{ or } 5.98 \text{ or } 5.99}$ $\text{Var} = \frac{(\text{Var}(b) \text{ or } \text{Var}(a))}{30.48^2 \text{ or } 929}$ <p>or</p> $\text{Var} = \left(\frac{(\text{Sd}(a) \text{ or } \text{Sd}(b))}{30.48} \right)^2$ $\text{Var}(n \text{ or } n-1) = \underline{0.0112 \text{ or } 0.0113 \text{ or } 0.0114 \text{ or } 0.0115}$	B1 Mdep1 A1	 3	CAO (5.98261) Dep on 9 < Var < 12 in (a) or (b) $(30.48^2 = 929.0304)$ Dep on 3 < Sd < 3.5 in (a) or (b) CAO $(0.0113 \text{ or } 0.0114)$
		Total	9	

Q	Solution	Marks	Total	Comments
4 (a)	$r = \frac{-0.524}{}$ $= \underline{-0.52 \text{ to } -0.53}$ $= \underline{-0.4 \text{ to } -0.6}$	B3 (B2) (B1)		AWRT AWFW AWFW (-0.52387)
	Attempt at $\sum x$ $\sum x^2$ $\sum y$ $\sum y^2$ & $\sum xy$ or Attempt at S_{xx} S_{yy} & S_{xy} Attempt at substitution into correct corresponding formula for r	(M1) (m1) (A1)		7631 4496183 5031 1966807 & 2943644 (all 5 attempted) 16786 19810 & -9553 (all 3 attempted) M0 mo A0 if r not in $[-1, +1]$ AWRT
	$r = \underline{-0.524}$		3	
(b)	Moderate or some negative (linear) correlation	Bdep1		Dependent on $-0.6 \leq r \leq -0.4$
Notes	1 Statements must include the words “moderate or some and negative” or “spurious” together with “correlation” or “association” or “relationship”; ignore additional comments unless clearly contradictory 2 Use of any of the following terms (even in conjunction with moderate or some): “strong or high or big or good or low or little or small or weak or slight or fairly or mild or medium or average or reasonable or pretty” \Rightarrow Bdep0			
	between total weight(s) of 4 apples and 3 bananas	B1		Context; providing $-1 < r < 1$
Notes	1 “As weights of (4) apples increase weights of (3) bananas decrease” (OE) Bdep0 B1 2 “As weights/ x increase weights/ y decrease” (OE) Bdep0 B0 3 Allow “mass(es) or gram(s)” instead of “weight(s)”			
SCs	Special cases 1 and 2 depend upon $-1 < r < 1$ 1 Any answer suggesting that the correlation between the two variables is nonsensical/tenuous/dodgy/rubbish/OE \Rightarrow B1 2 Any answer suggesting that the correlation between the weight(s) of 4 apples and 3 bananas is nonsensical/tenuous/dodgy/rubbish/OE \Rightarrow B2			
(c)	Howard’s claim suggests positive correlation but Hilda's data/calculated value shows negative correlation Disagree with or reject claim or Claim is (likely to be) false/incorrect/wrong/invalid/inaccurate/unsupported/ unreasonable	Bdep1 Bdep1		Correct comparison of claim and calculated value using positive and negative Dependent on $-1 < r < 0$ OE; dependent on Bdep1
			2	
		Total	7	

Q	Solution	Marks	Total	Comments
5(a)	Accept the equivalent percentage answers with %-sign (see GN5)			
(i)	$P(X < 105) = \underline{\mathbf{0.5 \text{ or } 1/2 \text{ or half or } 50\%}}$	B1	(1)	CAO; accept nothing else but ignore zeros after 0.5 (eg 0.50) Ignore additional words providing that they are not contradictory
(ii)	$P(X \neq 100) = \underline{\mathbf{1 \text{ or one or unity or } 100\%}}$	B1	(1)	CAO; accept nothing else but ignore zeros after decimal point (eg 1.00) Ignore additional words providing that they are not contradictory (eg certain so = 1)
(iii)	$P(X > 110) = P\left(Z > \frac{110 - 105}{4}\right) =$ $P(Z > 1.25) = 1 - 0.89435 = \underline{\mathbf{0.105 \text{ to } 0.106}}$	M1 A2	(3)	Standardising 110 with 105 and 4 but allow (105 – 110) AWFW (0.10565)
SCs	1 Answer of 0.894 to 0.895 \Rightarrow M1 A1 2 Correct seen standardisation with $0 < \text{incorrect answer} < 0.5 \Rightarrow$ M1 A1 3 Incorrect or no seen standardisation with $0 < \text{incorrect answer} < 0.5 \Rightarrow$ M1 A0			
(iv)	$P(102 < X < 108) = P(\underline{\mathbf{-0.75}} < Z < \underline{\mathbf{0.75}})$ $= (p - (1 - p)) \text{ or } (2p - 1)$ $= 0.77337 - (1 - 0.77337)$ $= \underline{\mathbf{0.546 \text{ to } 0.547}}$	M1 M1 A1	(3)	CAO -0.75 and +0.75 OE; $0 < p < 1$ Independent of previous M1 AWFW (0.54674)
			8	
(b)				
(i)	$5\% (0.05) \Rightarrow z = \underline{\mathbf{1.64 \text{ to } 1.65}}$ $\frac{\pm((150 \text{ or } 170) - 160)}{\sigma / s} = \pm(1.64 \text{ to } 1.65)$ $\sigma / s = \underline{\mathbf{6.06 \text{ or } 6.08 \text{ or } 6.10}}$	B1 M1 A1	3	AWFW (1.64485) Seen; ignore sign Standardising 150 with 160 and σ / s ; allow $(160 - (150 \text{ or } 170))$ and equating to $\pm(1.64 \text{ to } 1.65)$ and with consistent signs CAO (6.07957) Seen incorrect rounding \Rightarrow A0
Notes	1 Award max of B1 M0 A0 for any inconsistency of signs 2 Award of B0 \Rightarrow 0/3 marks here 3 Whilst the use of $z = 1.96$ ($\sigma = 5.1$) or $z = 1.28$ ($\sigma = 7.8$) \Rightarrow 0/3 marks here, a max of B1 M1 is available in (b)(ii)(B)			
	Parts (a) & (b)(i)	Total	11	

Q	Solution	Marks	Total	Comments
5	Continued			
	Parts (a) & (b)(i)	Total	11	
(b)(ii) (A)	$P(Y > 150) = 1 - 0.05 = 0.95$ $P(Y_1 \& Y_2 \& Y_3 > 150) = 0.95^3$ $= \underline{\mathbf{0.857 \text{ to } 0.858}}$	B1	(1)	AWFW (0.857375)
Note	1 A calculation of $P(Y > 150) = p$ followed by $p^3 \Rightarrow$ B1 only if result falls within above range			
(B)	$V(\bar{Y}) = \frac{(5.1 \text{ to } 7.9)^2}{3} \text{ or } \frac{(26 \text{ to } 61.1)}{3}$ <p style="text-align: center;">or 8.6 to 20.4</p> <p>or</p> $Sd(\bar{Y}) = \frac{5.1 \text{ to } 7.9}{\sqrt{3}} \text{ or } \underline{\mathbf{2.93 \text{ to } 4.57}}$	B1		AWFW (12.32038) Can be implied by what follows AWFW (3.51004)
	$P(\bar{Y} > 162.5) = P\left(Z > \frac{162.5 - 160}{(5.1 \text{ to } 7.9)/\sqrt{3}}\right)$ $= P(Z > \underline{\mathbf{0.70 \text{ to } 0.72}})$ $= \underline{\mathbf{0.235 \text{ to } 0.242}}$	M1 A1 A1		Standardising 162.5 with 160 and (5.1 to 7.9)/√3 (OE) ; allow (160 – 162.5) AWFW; ignore sign (0.71224) AWFW (0.23816)
			(4) 5	
Notes	1 Do not give BOD for unclear/dubious/questionable identifications of (A) & (B) 2 If answers to (A) & (B) are not identified, then mark as (A) followed by (B) 3 If answers to (A) & (B) are switched, then 0/5 marks 4 In (B), award of B0 \Rightarrow 0/4 marks 5 In (B), cubing (0.235 to 0.242) \Rightarrow B1 M1 A1 A0 (ie not ISW) 6 In (B), for information, use of (5.1 to 7.9) $\Rightarrow z = (0.548 \text{ to } 0.849) \Rightarrow p = (0.197 \text{ to } 0.292)$			
SC	1 Use of distribution of total in (B): B1 for $Sd = (5.1 \text{ to } 7.9)\sqrt{3}$ (OE); M1 for $P(Z > (487.5 - 480)/((5.1 \text{ to } 7.9)\sqrt{3}))$ (OE); A1 for (0.70 to 0.72) A1 for 0.235 to 0.242 (AWFW); award of B0 \Rightarrow 0/4 marks			
		Total	16	

Q	Solution	Marks	Total	Comments
6 (a) (i)	Accept the equivalent percentage answers with %-sign (see GN5)			
	$P(A_1 \cap A_2 \cap A_3) = 0.85^3$ $= \underline{\underline{0.614}}$	B1	(1)	AWRT (0.614125)
(ii)	$P(A_1 \cap A_2 \cap B) = 0.85^2 \times 0.10$ or (0.0722 to 0.0723) or 289/4000 $\times \underline{\underline{3}}$ $= \underline{\underline{0.216 \text{ to } 0.217}}$	M1 A1 A1		(3) OE; do not accept additional terms (0.07225) OE AWFW (0.21675)
(iii)	$P(A \cap B \cap C) = 0.85 \times 0.10 \times 0.05$ or (0.0042 to 0.0043) or 17/4000 $\times \underline{\underline{6}}$ $= \underline{\underline{0.025 \text{ to } 0.026}}$	M1 A1 A1	(3) OE; do not accept additional terms (0.00425) OE AWFW (0.0255)	
Notes	1 Deduct 1 mark for at least one fractional final answer of (i) 4913/8000; (ii) 867/4000; (iii) 51/2000 2 Award A0 for each alternative fractional final answer [eg (ii) 4335/20000]			
	Part (a)	Total	7	

Q	Solution	Marks	Total	Comments																							
6	Accept the equivalent percentage answers with %-sign (see GN5)																										
	Part(a)	Total	7																								
(b)																											
(i)	$P(OD') = 0.10 \times 0.75 + 0.05 \times 0.10$ <p style="text-align: center;">PLUS $0.85 (\times 1)$</p> $= 0.85 + 0.075 + 0.005 = \underline{\underline{0.93}}$	M1 A1 A1		CAO CAO																							
	or																										
	$P(OD) = 0.10 \times 0.25 + 0.05 \times 0.90$ $+ 0.85 \times 0$ $= 0.025 + 0.045 = \underline{\underline{0.07}}$ $P(OD') = 1 - 0.07 = \underline{\underline{0.93}}$	(M1) (A1) (A1)		See SC 3 below CAO CAO																							
	or																										
	(y) <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">Side Effect</th> <th rowspan="2">Total</th> </tr> <tr> <th>None</th> <th>Slight</th> <th>Severe</th> </tr> </thead> <tbody> <tr> <th>Change</th> <td>0</td> <td>2.5</td> <td>4.5</td> <td>7.0</td> </tr> <tr> <th>No change</th> <td>85</td> <td>7.5</td> <td>0.5</td> <td>93.0</td> </tr> <tr> <th>Total</th> <td>85</td> <td>10.0</td> <td>5.0</td> <td>100.0</td> </tr> </tbody> </table>		Side Effect			Total	None	Slight	Severe	Change	0	2.5	4.5	7.0	No change	85	7.5	0.5	93.0	Total	85	10.0	5.0	100.0	(B2) (B1)		Accept probabilities rather than percentages (0, 2.5, 4.5) or (85, 7.5, 0.5) CAO 0.93 CAO
	Side Effect			Total																							
	None	Slight	Severe																								
Change	0	2.5	4.5	7.0																							
No change	85	7.5	0.5	93.0																							
Total	85	10.0	5.0	100.0																							
			(3)																								
Note	1 Accept fractional answers of 93/100 and 7/100																										
SCs	1 In (α) , $P(OD') = 0.10 \times 0.75 + 0.05 \times 0.10 + 0.15 (\times 1) = 0.23 \Rightarrow$ M1 A0 A0 (max 1 mark) 2 In (β) , $P(OD) = 0.10 \times 0.25 + 0.05 \times 0.90 = 0.07 \Rightarrow$ M1 A1 then $P(OD') = 0.93 \Rightarrow$ A1 (max 3 marks) 3 In (β) , $P(OD) = 0.10 \times 0.25 + 0.05 \times 0.90 + (\text{not } 0.85) \times (0) = 0.07 \Rightarrow$ M1 A0 then $P(OD') = 0.93 \Rightarrow$ A1 (max 2 marks) 4 In (β) , $P(OD) = 0.10 \times 0.25 + 0.05 \times 0.90 + (\text{non-zero term}) \Rightarrow$ M1 A0 A0 (max 1 mark)																										
(ii)	$P(OD B \cup C) = \frac{1-0.93}{0.10+0.05} \text{ or } \frac{0.07}{0.10+0.05}$ $= \underline{\underline{7/15 \text{ or } 0.466 \text{ to } 0.467 \text{ or } 0.4\dot{6}}}$	M1 M1 A1		Numerator; OE Denominator (See Notes 1 & 2 below) CAO/AWFW/CAO (0.46667)																							
	or																										
	$P(OD B \cup C) = \frac{2}{3} \times 0.25 + \frac{1}{3} \times 0.9$ $= \underline{\underline{7/15 \text{ or } 0.466 \text{ to } 0.467 \text{ or } 0.4\dot{6}}}$	(M1) (M1) (A1)		Either term (OE) PLUS other term (OE) CAO/AWFW/CAO (0.46667)																							
	or																										
	From table, $P(OD B \cup C)$																										
	$= \underline{\underline{7/15 \text{ or } 0.466 \text{ to } 0.467 \text{ or } 0.4\dot{6}}}$	(B3)		CAO/AWFW/CAO (0.46667)																							
			(3)																								
Notes	1 A mark of M1 may be available in a fraction even if the resultant probability answer is greater than 1 2 Values of $(1 - 0.93)$ or 0.07 or 0.15 seen but not in a fraction and with no correct answer \Rightarrow M0 M0 (A0)																										
			6																								
		Total	13																								

Q	Solution	Marks	Total	Comments
7	Accept 3 dp rounding of probabilities from tables	Accept the equivalent percentage answers with %-sign (see GN5)		
(a)				
(i)	$P(\text{Blond} = 5) = \binom{30}{5} (0.25)^5 (0.75)^{25}$ $= 142506 \times 0.00097656 \times 0.00075254$ or $= 0.2026 - 0.0979$ $= \underline{\underline{0.104 \text{ to } 0.105}}$	M1 A1	2	Correct expression Can be implied by a correct answer Ignore additional expressions AFWW (0.104728 / 0.1047)
(ii)	$P(\text{Blond} < 10) = \underline{\underline{0.803}}$	B1	1	AWRT (0.8034)
(iii)	$P(6 \leq \text{Blond} \leq 12) =$ $0.9784 \text{ or } 0.9493 \quad (p_1)$ MINUS $0.2026 \text{ or } 0.3481 \quad (p_2)$ $= \underline{\underline{0.775 \text{ to } 0.776}}$	M1 M1 A1	3	Seen as first term in a subtraction Seen as second term in a subtraction AFWW (0.7758)
Notes	1 For no method or calculation of individual terms: award B3 for 0.775 to 0.776 (AWFW); B2 for 0.746 to 0.747 (AWFW); B2 for 0.630 to 0.631 (AWFW); B2 for 0.601 to 0.602 (AWFW); B0 for anything else 2 Answers seen using [(0.7974 or 0.6519) – (0.0216 or 0.0507)] [ie (1 – p ₂) – (1 – p ₁)] ⇒ M1 M1 A1 max 3 Answers seen using [1 – (p ₁ – p ₂)] even after (p ₁ – p ₂) [eg 1 – (0.9784 – 0.2026) = 0.2242] ⇒ 0 marks 4 Use of p ₁ × p ₂ or p ₁ ÷ p ₂ or p ₁ + p ₂ or p ₁ only or p ₂ only ⇒ 0 marks			
(iv)	$\text{Mean} = np = 7.5 \Rightarrow P(\text{Blond} \geq 8)$ $= 1 - 0.5143$	M2		
	$= (1 - 0.6736) \text{ or } 0.3264$			
	or			
	$= 0.5143$	(M1)		
	or			
	$= (1 - 0.3481) \text{ or } 0.6519$			
	$= \underline{\underline{0.485 \text{ to } 0.486}}$	A1	3	AWFW (0.4857)
Note	1 For calculation of individual terms or no method: award B3 for 0.485 to 0.486 (AWFW); B1 for 0.326 to 0.327 (AWFW); B1 for 0.514 to 0.515 (AWFW); B1 for 0.651 to 0.652 (AWFW); B0 for anything else			
	Part(a)	Total	9	

Q	Solution	Marks	Total	Comments
7	Accept 3 dp rounding of probabilities from tables	Accept the equivalent percentage answers with %-sign (see GN5)		
	Part(a)	Total	9	
(b) (i)	Mean = $np = 16$	B1		Equating; seen or used
	$np(1-p)$ or npq or $\sqrt{np(1-p)}$ or \sqrt{npq} $= 2.4^2$ or 5.76 or 2.4 but not $\sqrt{2.4}$	M1		Equating; seen or used
	$np(1-p)$ or $npq = 2.4^2$ or 5.76 $p = \underline{0.64}$ and $n = \underline{25}$	A1 A1 A1		Equating; seen or used Each CAO
Notes	1 Equating npq to 2.4 (OE) then \Rightarrow B1 M1 A0 A0 A0 (max) followed by M0 A0 in (ii) 2 For any method, answer of $p = 0.64$ (CAO) and $n = 25$ (CAO) \Rightarrow 5 marks 3 For method of 'trial & improvement': B1 (equating/use of $np = 16$); M1 (at least one seen trial combination of either integer n or $0 < p < 1$); m1 (at least one seen attempt at evaluating npq with both integer n and $0 < p < 1$ but comparison with 5.76/2.4 not required); A1 ($p = 0.64$ CAO); A1 ($n = 25$ CAO)			
(ii)	$P(Y = 20) = \binom{25}{20} (0.64)^{20} (0.36)^5$ $= 53130 \times 0.00013292 \times 0.0060466$ $= \underline{0.0426 \text{ to } 0.0428}$	M1 A1		Correct expression Can be implied by a correct answer Do not ignore additional expressions AFWW (0.042702)
		Total	16	