



Surname _____

Other Names _____

Centre Number _____

Candidate Number _____

Candidate Signature _____

AS

CHEMISTRY

Paper 2 Organic and Physical Chemistry

7404/2

Thursday 23 May 2019 Morning

Time allowed: 1 hour 30 minutes

For this paper you must have:

- **the Periodic Table/Data Sheet, provided as an insert (enclosed)**
- **a ruler with millimetre measurements**
- **a scientific calculator, which you are expected to use where appropriate.**

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.

[Turn over]



BLANK PAGE



INSTRUCTIONS

- **Use black ink or black ball-point pen.**
- **Answer ALL questions.**
- **You must answer the questions in the spaces provided. Do NOT write on blank pages.**
- **All working must be shown.**
- **Do all rough work in this book. Cross through any work you do not want to be marked.**

INFORMATION

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 80.**

ADVICE

You are advised to spend about 65 minutes on SECTION A and 25 minutes on SECTION B.

DO NOT TURN OVER UNTIL TOLD TO DO SO




SECTION A


Answer ALL questions in this section.

0	1
---	---


The structures of three organic compounds A, B and C are shown.



Compound A



Compound B



Compound C

These compounds can be distinguished by simple test-tube reactions.



For each pair of compounds in questions 01.1 and 01.2, give a reagent (or combination of reagents) that could be added separately to each compound to distinguish between them.

State what is observed in each case.

0 1 . 1 Compounds A and B [3 marks]

Reagent _____

Observation with A _____

Observation with B _____

[Turn over]



BLANK PAGE



0 1 . 2 Compounds A and C [3 marks]

Reagent _____

Observation with A _____

Observation with C _____


6

[Turn over]



0 2

Bromoethane reacts with potassium cyanide to form compound D.



Compound D

0 2 . 1

**Outline the mechanism for this reaction.
[2 marks]**

0 2 . 2

Give the IUPAC name of D. [1 mark]



0 2 . 3 Calculate the percentage atom economy for the formation of D in this reaction.

Give your answer to the appropriate number of significant figures. [2 marks]

% atom economy _____

5

[Turn over]



0 3 This question is about enthalpy changes.

0 3 . 1 A student determined the enthalpy of combustion of cyclohexane (C_6H_{12}).

The student

- placed a pure sample of cyclohexane in a spirit burner
- placed the spirit burner under a beaker containing 50.0 g of water and ignited the cyclohexane
- extinguished the flame after a few minutes.

The results for the experiment are shown in TABLE 1.

TABLE 1

Initial temperature of the water / °C	19.1
Initial mass of spirit burner and cyclohexane / g	192.730
Final mass of spirit burner and cyclohexane / g	192.100

The student determined from this experiment that the enthalpy of combustion of cyclohexane is $-1216 \text{ kJ mol}^{-1}$



Use the data to calculate the final temperature of the water in this experiment.

The specific heat capacity of water = $4.18 \text{ J K}^{-1} \text{ g}^{-1}$

The relative molecular mass (M_r) of cyclohexane = 84.0

[4 marks]

Final temperature of the water _____ °C

[Turn over]



0 3 . 2 A data book value for the enthalpy of combustion of cyclohexane is $-3920 \text{ kJ mol}^{-1}$

The student concluded that the temperature rise recorded in the experiment was smaller than it should have been.

Suggest a practical reason for this. [1 mark]

0 3 . 3 TABLE 2 gives some values of standard enthalpies of combustion ($\Delta_c H^\theta$).

TABLE 2

Substance	C(s)	H ₂ (g)	C ₆ H ₁₂ (l)
Standard enthalpy of combustion, $\Delta_c H^\theta / \text{kJ mol}^{-1}$	-394	-286	-3920



Use the data in TABLE 2 to calculate the enthalpy change for the reaction represented by this equation



[3 marks]

Enthalpy change _____ kJ mol⁻¹

[Turn over]

8



0 4

This question is about fossil fuels.

0 4 . 1

The petrol fraction from crude oil contains octane (C_8H_{18}).

Give an equation for the complete combustion of octane. [1 mark]

0 4 . 2

The combustion of petrol in car engines produces the pollutant nitrogen monoxide.

Give an equation for a reaction that removes nitrogen monoxide in a catalytic converter. [1 mark]




BLANK PAGE

[Turn over]



0 4 . 3 Sulfur dioxide is produced in the combustion of fossil fuels. The total emissions of sulfur dioxide in the UK have fallen dramatically since 1970.

Sulfur dioxide is now removed from the flue gases in power stations by reaction with calcium oxide.



In 1970, the total UK emissions of sulfur dioxide were 6.49 million tonnes (1 tonne = 1000 kg).

Calculate the mass, in kilograms, of calcium oxide needed to react with this mass of sulfur dioxide.

Give your answer in standard form.
[2 marks]



Mass of calcium oxide _____ kg

4

[Turn over]



0 5

Methanol (CH₃OH) is an important alcohol with many uses.

0 5 . 1

Draw a diagram to show how two methanol molecules interact with each other through hydrogen bonding in the liquid phase.

Include all partial charges and all lone pairs of electrons in your diagram. [3 marks]



0 5 . 2

The bond angle around the oxygen atom in methanol is slightly smaller than the regular tetrahedral angle of 109.5°

Explain why this bond angle is smaller than 109.5° [1 mark]

[Turn over]



0 5 . 3

Methanol is made by the reaction of carbon monoxide with hydrogen.



The reaction uses a copper-based catalyst, a pressure of 10 MPa and a temperature of 550 K

These conditions are used to provide a balance between equilibrium yield, reaction rate and cost.


Describe how the use of a catalyst, and changes in pressure and temperature, each affect equilibrium yield, reaction rate and cost. [6 marks]



0 6

Propene reacts with concentrated sulfuric acid to form two isomers, E and F.

The structure of E is shown.

**0 6 . 1**

Name and outline the mechanism for the formation of E in this reaction. [5 marks]

Name of mechanism

Mechanism



0 6 . 2 Draw the structure of F. [1 mark]

[Turn over]



06.3

Explain why more of isomer E than isomer F is formed in this reaction. [2 marks]

8



BLANK PAGE

[Turn over]



0 7

Propanedioic acid contains two carboxylic acid groups. It is a solid organic acid that is soluble in water.

0 7 . 1

Draw the skeletal formula of propanedioic acid. [1 mark]

0 7 . 2

Describe how to prepare 250 cm³ of an aqueous standard solution of propanedioic acid containing an accurately measured mass of the acid. Include essential practical details in your answer. [6 marks]



07.3

Calculate the mass, in mg, of propanedioic acid ($M_r = 104.0$) needed to prepare 250 cm^3 of a $0.00500 \text{ mol dm}^{-3}$ solution.
[2 marks]

Mass of propanedioic acid _____ mg

9



BLANK PAGE


[Turn over]



0 8

Propanal can be prepared by the oxidation of propan-1-ol with acidified potassium dichromate(VI).

An ionic equation for this reaction is

**0 8 . 1**

Calculate the minimum volume, in cm^3 , of 0.40 mol dm^{-3} potassium dichromate(VI) solution needed to oxidise 6.0 cm^3 of propan-1-ol to propanal.

M_r of propan-1-ol = 60.0

Density of propan-1-ol = 0.80 g cm^{-3}
[3 marks]



Minimum volume _____ cm³


[Turn over]



0 8 . 2 The reaction is done in a pear-shaped flask.


Complete the diagram to show the assembled apparatus needed to prepare propanal from propan-1-ol in this way.

Label the diagram. [3 marks]



0 9

The compound 1,2-dichlorotetrafluoroethane is a CFC that was previously used in refrigerators as a coolant.

**0 9 . 1**

Molecules of 1,2-dichlorotetrafluoroethane can break down in the upper atmosphere to form chlorine radicals.

Give an equation to show the breakdown of one molecule of 1,2-dichlorotetrafluoroethane to form one chlorine radical and one other species. [1 mark]

0 9 . 2

Give TWO equations to show how chlorine radicals catalyse the decomposition of ozone. [2 marks]

[Turn over]



09.3

Butane can be used as a replacement for CFCs in refrigerators.

During its use, the butane is repeatedly converted from liquid to gas and then back to liquid. Liquid butane expands as it turns into a gas.

- Calculate the volume, in cm^3 , of 38.8 g of butane gas at 272 K and 101 kPa
(the gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$)
(M_r of butane = 58.0)
- Calculate the volume, in cm^3 , of 38.8 g of liquid butane.
(density of liquid butane = 0.60 g cm^{-3})
- Use your answers to calculate the factor by which butane expands in volume when it changes from a liquid to a gas.

Show your working. [6 marks]



Volume of butane gas _____ cm^3

Volume of liquid butane _____ cm^3

Expansion factor _____

[Turn over]

9




SECTION B




Answer ALL questions in this section.

Only ONE answer per question is allowed.
For each answer completely fill in the circle alongside
the appropriate answer.


CORRECT METHOD




WRONG METHODS



If you want to change your answer you must
cross out your original answer as shown.



If you wish to return to an answer previously
crossed out, ring the answer you now wish to
select as shown.



You may do your working in the blank space around
each question but this will not be marked.
Do NOT use additional sheets for this working.



1	0
---	---

A 'drink-driving' offence is committed if the blood alcohol level of a driver is over 80 mg of ethanol per 100 cm³ of blood. What is the concentration, in mol dm⁻³, of ethanol if there are 80 mg of ethanol ($M_r = 46.0$) per 100 cm³ of blood?
[1 mark]

A 0.00017

B 0.0017

C 0.017

D 1.7

[Turn over]



1	1
---	---

Which statement is correct for the distribution curve of molecular energies in a gas? [1 mark]

A The curve is symmetrical about the maximum.

B There are always some molecules with zero energy.


C The position of the maximum of the curve is not dependent on the temperature.

D The mean energy of the molecules is greater than the most probable energy of the molecules.



1	2
---	---

When one mole of ammonia is heated to a given temperature, 50 % of it dissociates and the following equilibrium is established.



What is the total amount, in moles, of gas in this equilibrium mixture? [1 mark]

A 1.5

B 2.0



C 2.5

D 3.0


[Turn over]




1 3 Which compound is NOT an isomer of the following compound? [1 mark]




A $\text{CH}_3\text{CH}_2\text{COCH}_3$



B $\text{CH}_3\text{CH}=\text{CHCH}_2\text{OH}$




C $(\text{CH}_3)_2\text{CHCHO}$




D $\text{CH}_2=\text{CHCH}_2\text{CHO}$




1 4 How many isomers are there of C_3H_9N ?
[1 mark]




A 2



B 3



C 4





D 5

[Turn over]



1 5 Which equation represents a propagation step? [1 mark]



1 6 Which compound can react with ammonia to produce propylamine? [1 mark]

A $\text{CH}_3\text{CH}=\text{CH}_2$

B $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$

C $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$

D $\text{CH}_3\text{CH}_2\text{CH}_3$

[Turn over]



1 7

Which statement is NOT correct about $\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}_2\text{Br}$? [1 mark]

A It displays *E-Z* isomerism.

B It forms an addition polymer.

C It reacts with electrophiles.

D It decolourises bromine water.

1 8

Which compound can be oxidised to form $(\text{CH}_3)_2\text{CHCOCH}_3$? [1 mark]

A 2-methylpropan-1-ol

B 2,2-dimethylpropanol

C 2-methylbutan-2-ol

D 3-methylbutan-2-ol

[Turn over]



1 9

Which species can act as a nucleophile?
[1 mark]

A NH_4^+

B CH_3OH

C CH_4

D H^+



2 0 Which alcohol forms a mixture of alkenes when dehydrated? [1 mark]

A propan-1-ol

B propan-2-ol

C pentan-1-ol

D pentan-2-ol

[Turn over]



2 1 Which compound has the highest boiling point? [1 mark]

A $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$

B $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$

C $\text{CH}_3\text{CH}_2\text{CHO}$

D $\text{CH}_3\text{CH}_2\text{COOH}$



2	2
---	---

Which compound could NOT be produced by reacting 2-bromo-3-methylbutane with sodium hydroxide? [1 mark]

A 2-methylbut-1-ene

B 3-methylbut-1-ene

C 2-methylbut-2-ene

D 3-methylbutan-2-ol


[Turn over]



2 3

The infrared spectrum of an organic compound is shown.

Transmittance
/ %



Which compound produces this spectrum?
[1 mark]

A ethanoic acid

B 4-hydroxybutanone

C propan-1-ol

D prop-2-en-1-ol



- 2 4** The heat released when 1.00 g of ethanol ($M_r = 46.0$) undergoes complete combustion is 29.8 kJ

What is the heat released by each molecule, in joules, when ethanol undergoes complete combustion?

(the Avogadro constant $L = 6.022 \times 10^{23} \text{ mol}^{-1}$)
[1 mark]

A $2.28 \times 10^{-18} \text{ J}$

B $4.95 \times 10^{-20} \text{ J}$

C $2.28 \times 10^{-21} \text{ J}$

D $4.95 \times 10^{-23} \text{ J}$

15

END OF QUESTIONS



BLANK PAGE

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
Section B	
TOTAL	

Copyright Information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.

IB/M/CH/Jun19/7404/2/E3