



Surname _____

Other Names _____

Centre Number _____

Candidate Number _____

Candidate Signature _____

A-level

CHEMISTRY

Paper 1 Inorganic and Physical Chemistry

7405/1

Tuesday 4 June 2019 Afternoon

Time allowed: 2 hours

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.

[Turn over]



For this paper you must have:

- **the Periodic Table/Data Sheet, provided as an insert (enclosed)**
- **a ruler with millimetre measurements**
- **a scientific calculator, which you are expected to use where appropriate.**

INSTRUCTIONS

- **Use black ink or black ball-point pen.**
- **Answer ALL questions.**
- **You must answer the questions in the spaces provided. Do NOT write on blank pages.**
- **All working must be shown.**
- **Do all rough work in this book. Cross through any work you do not want to be marked.**



INFORMATION

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 105.**

**DO NOT TURN OVER UNTIL TOLD TO
DO SO**



Answer ALL questions in the spaces provided.

0 1

FIGURE 1 shows an incomplete Born-Haber cycle for the formation of caesium iodide. The diagram is not to scale.

FIGURE 1




TABLE 1 gives values of some standard enthalpy changes.

TABLE 1


| Name of enthalpy change | $\Delta H^\ominus / \text{kJ mol}^{-1}$ |
|--|---|
| Enthalpy of atomisation of caesium | +79 |
| First ionisation energy of caesium | +376 |
| Electron affinity of iodine | -314 |
| Enthalpy of lattice formation of caesium iodide | -585 |
| Enthalpy of formation of caesium iodide | -337 |

[Turn over]



01.1

Complete FIGURE 1, on page 4, by writing the formulas, including state symbols, of the appropriate species on each of the two blank lines. [2 marks]



0 1 . 2

Use FIGURE 1 and the data in TABLE 1, on page 5, to calculate the standard enthalpy of atomisation of iodine. [2 marks]

7


Standard enthalpy of atomisation of iodine

_____ kJ mol⁻¹

[Turn over]



BLANK PAGE



0 1 . 3

The enthalpy of lattice formation for caesium iodide in TABLE 1, on page 5, is a value obtained by experiment.

The value obtained by calculation using the perfect ionic model is -582 kJ mol^{-1}


Deduce what these values indicate about the bonding in caesium iodide. [1 mark]

[Turn over]



01.4

Use data from TABLE 2 to show, in the space on the opposite page, that this reaction is NOT feasible at 298 K



$$\Delta H^\ominus = +337 \text{ kJ mol}^{-1}$$

TABLE 2

| | CsI (s) | Cs (s) | I ₂ (s) |
|--|---------|--------|--------------------|
| $S^\ominus / \text{J K}^{-1} \text{ mol}^{-1}$ | 130 | 82.8 | 117 |

[4 marks]



[Turn over]

9



0 2


Time of flight (TOF) mass spectrometry can be used to analyse large molecules such as the pentapeptide, leucine enkephalin (P).

P is ionised by electrospray ionisation and its mass spectrum is shown in FIGURE 2.

12



FIGURE 2



[Turn over]




02.1

Describe the process of electrospray ionisation.

Give an equation to represent the ionisation of P in this process. [4 marks]

Description

14



Equation

[Turn over]

15



BLANK PAGE



02.2

What is the relative molecular mass of P?


Tick (✓) ONE box. [1 mark]

555

556

557

[Turn over]



0 2 . 3

A molecule Q is ionised by electron impact in a TOF mass spectrometer.

The Q^+ ion has a kinetic energy of $2.09 \times 10^{-15} \text{ J}$

This ion takes $1.23 \times 10^{-5} \text{ s}$ to reach the detector.

The length of the flight tube is 1.50 m

Calculate the relative molecular mass of Q.

$$KE = \frac{1}{2} mv^2 \quad \text{where } m = \text{mass (kg) and}$$
$$v = \text{speed (m s}^{-1}\text{)}$$

The Avogadro constant,

$$L = 6.022 \times 10^{23} \text{ mol}^{-1} \text{ [5 marks]}$$



Relative molecular mass

[Turn over]

| |
|-----------|
| |
| 10 |



| | |
|---|---|
| 0 | 3 |
|---|---|

This question is about periodicity, the Period 4 elements and their compounds.

| | | | |
|---|---|---|---|
| 0 | 3 | . | 1 |
|---|---|---|---|

**State the meaning of the term periodicity.
[1 mark]**

| | | | |
|---|---|---|---|
| 0 | 3 | . | 2 |
|---|---|---|---|

Identify the element in Period 4 with the highest electronegativity value. [1 mark]



0 3 . 3

Identify the element in Period 4 with the largest atomic radius.

Explain your answer. [3 marks]

Element _____


Explanation _____

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 3 | . | 4 |
|---|---|---|---|


The equations for two reactions of arsenic(III) oxide are shown.



Name the property of arsenic(III) oxide that describes its ability to react in these two ways. [1 mark]

| | | | |
|---|---|---|---|
| 0 | 3 | . | 5 |
|---|---|---|---|


Complete the equation for the formation of arsenic hydride. [1 mark]



[Turn over]

| |
|---|
| |
| 7 |

FIGURE 3



BLANK PAGE

[Turn over]



0 4

FIGURE 3, on page 24, shows some reactions of aqueous iron ions.

0 4 . 1

Give the formula of PRECIPITATE J and state its colour.

26

Give an equation for REACTION 1. [3 marks]

Formula of J _____




Colour

Equation

[Turn over]



Repeat of FIGURE 3



04.2

**Give the formula of L and an equation for REACTION 2.
[2 marks]**

Formula of L _____


29

Equation _____

[Turn over]



Repeat of FIGURE 3



0 4 . 3

Suggest a reagent for REACTION 3. [1 mark]

0 4 . 4

**Give the formula of PRECIPITATE M and state its colour.
[2 marks]**

Formula of M _____

Colour _____

[Turn over]



04.5

Transition metal complexes have different shapes and many show isomerism.

Describe the different shapes of complexes and show how they lead to different types of isomerism.

Use examples of complexes of cobalt(II) and platinum(II).

You should draw the structures of the examples chosen. [6 marks]

BLANK PAGE

[Turn over]



| | |
|---|---|
| 0 | 5 |
|---|---|

This question is about some Group 7 compounds.

| | | | |
|---|---|---|---|
| 0 | 5 | . | 1 |
|---|---|---|---|

Solid sodium chloride reacts with concentrated sulfuric acid.

**Give an equation for this reaction.
State the role of the sulfuric acid in this reaction. [2 marks]**

Equation

Role



05.2

Fumes of sulfur dioxide are formed when sodium bromide reacts with concentrated sulfuric acid.

For THIS reaction

- **give an equation**
- **give ONE other observation**
- **state the role of the sulfuric acid.**

[3 marks]

Equation

Observation

Role

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 5 | . | 3 |
|---|---|---|---|

Chlorine reacts with hot aqueous sodium hydroxide as shown in the equation.



Give the oxidation state of chlorine in NaClO_3 and in NaCl [1 mark]

NaClO_3 _____

NaCl _____

05.4

**State, in terms of redox, what happens to chlorine in the reaction in Question 05.3.
[1 mark]**

[Turn over]

05.5

Solution Y contains TWO different negative ions.

To a sample of solution Y in a test tube a student adds

- **silver nitrate solution**
- **then an excess of dilute nitric acid**
- **finally an excess of concentrated ammonia solution.**

The observations after each addition are recorded in TABLE 3.

TABLE 3

| REAGENT ADDED TO SOLUTION Y | OBSERVATION |
|---|---|
| silver nitrate solution | cream precipitate containing compound D and compound E |
| excess dilute nitric acid | cream precipitate D and bubbles of gas F |
| excess concentrated ammonia solution | colourless solution containing complex ion G |

[Turn over]



BLANK PAGE



Give the formulas of D, E and F.

Give an IONIC equation to show the formation of E.

Give an equation to show the conversion of D into G. [6 marks]

Formula of D _____

Formula of E _____

Formula of F _____

Ionic equation to form E

Equation to show the conversion of D into G

[Turn over]



| | |
|---|---|
| 0 | 6 |
|---|---|



A student does an experiment to determine the percentage of copper in an alloy.

The student

- reacts 985 mg of the alloy with concentrated nitric acid to form a solution (all of the copper in the alloy reacts to form aqueous copper(II) ions)**
- pours the solution into a volumetric flask and makes the volume up to 250 cm³ with distilled water**
- shakes the flask thoroughly**
- transfers 25.0 cm³ of the solution into a conical flask and adds an excess of potassium iodide**
- uses exactly 9.00 cm³ of 0.0800 mol dm⁻³ sodium thiosulfate (Na₂S₂O₃) solution to react with all the iodine produced.**



The equations for the reactions are



[Turn over]



| | | | |
|---|---|---|---|
| 0 | 6 | . | 1 |
|---|---|---|---|

Calculate the percentage of copper by mass in the alloy.

Give your answer to the appropriate number of significant figures. [6 marks]

% copper _____

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 6 | . | 2 |
|---|---|---|---|

Suggest TWO ways that the student could reduce the percentage uncertainty in the measurement of the volume of sodium thiosulfate solution, using the same apparatus as this experiment.
[2 marks]

1 _____

2 _____



| | | | |
|---|---|---|---|
| 0 | 6 | . | 3 |
|---|---|---|---|

State the role of iodine in the reaction with sodium thiosulfate. [1 mark]

| | | | |
|---|---|---|---|
| 0 | 6 | . | 4 |
|---|---|---|---|

Give the full electron configuration of a copper(II) ion. [1 mark]

[Turn over]

BLANK PAGE

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 6 | . | 6 |
|---|---|---|---|

Iodine vaporises easily.

Calculate the volume, in cm^3 , that 5.00 g of iodine vapour occupies at 185 °C and 100 kPa

The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

**Give your answer to 3 significant figures.
[4 marks]**

Volume _____ cm^3

[Turn over]

| |
|----|
| |
| 16 |



| | |
|---|---|
| 0 | 7 |
|---|---|

Sulfur trioxide decomposes on heating to form an equilibrium mixture containing sulfur dioxide and oxygen.



| | | | |
|---|---|---|---|
| 0 | 7 | . | 1 |
|---|---|---|---|

A sample of sulfur trioxide was heated and allowed to reach equilibrium at a given temperature.

The equilibrium mixture contained 6.08 g of sulfur dioxide.

Calculate the mass, in g, of oxygen gas in the equilibrium mixture. [2 marks]

Mass _____ g

[Turn over]



07.2

A different mass of sulfur trioxide was heated and allowed to reach equilibrium at 1050 K



The amounts of each substance in the equilibrium mixture are shown in TABLE 4.

TABLE 4

| Substance | Amount at equilibrium / mol |
|------------------------|------------------------------------|
| sulfur trioxide | 0.320 |
| sulfur dioxide | 1.20 |
| oxygen | 0.600 |

BLANK PAGE

[Turn over]



60

For this reaction at 1050 K the equilibrium constant, $K_p = 7.62 \times 10^5 \text{ Pa}$

Calculate the mole fraction of each substance at equilibrium.

Give the expression for the equilibrium constant, K_p

Calculate the total pressure, in Pa, of this equilibrium mixture. [4 marks]



Mole fraction SO_3

Mole fraction SO_2

Mole fraction O_2

K_p

Total pressure _____ Pa

[Turn over]



BLANK PAGE


[Turn over]



07.4

Use data from Question 07.3 to calculate the value of K_p , at 500 K, for the equilibrium represented by this equation.

Deduce the units of K_p



[2 marks]

K_p

Units

[Turn over]

| |
|----|
| |
| 10 |



| | |
|---|---|
| 0 | 8 |
|---|---|

This question is about structure and bonding.

| | | | |
|---|---|---|---|
| 0 | 8 | . | 1 |
|---|---|---|---|

Draw a diagram on the opposite page to show the strongest type of interaction between two molecules of ethanol ($\text{C}_2\text{H}_5\text{OH}$) in the liquid phase.

Include all lone pairs and partial charges in your diagram. [3 marks]

[Turn over]



08.2

Methoxymethane (CH_3OCH_3) is an isomer of ethanol.

TABLE 5 shows the boiling points of ethanol and methoxymethane.

TABLE 5

| Compound | Boiling point / °C |
|-----------------------|---------------------------|
| ethanol | 78 |
| methoxymethane | -24 |

In terms of the intermolecular forces involved, explain the difference in boiling points. [3 marks]

| | | | |
|---|---|---|---|
| 0 | 8 | . | 3 |
|---|---|---|---|

Draw the shape of the POCl_3 molecule and the shape of the ClF_4^- ion.

Include any lone pairs of electrons that influence the shapes.

In a POCl_3 molecule the oxygen atom is attached to the phosphorus atom by a double bond that uses two electrons from phosphorus.

Name each shape.

Suggest a value for the bond angle in ClF_4^-

[5 marks]



Shape of POCl_3

Shape of ClF_4^-

[Turn over]



BLANK PAGE



73

Name of shape of POCl_3

Name of shape of ClF_4^-

Bond angle in ClF_4^-

[Turn over]

| |
|----|
| |
| 11 |



| | |
|---|---|
| 0 | 9 |
|---|---|

This question is about different pH values.

| | | | |
|---|---|---|---|
| 0 | 9 | . | 1 |
|---|---|---|---|

**For pure water at 40 °C, pH = 6.67
A student thought that the water was acidic.**

Explain why the student was incorrect.

Determine the value of K_w at this temperature. [4 marks]


Explanation _____

K_w _____ $\text{mol}^2 \text{dm}^{-6}$

[Turn over]



FIGURE 4



BLANK PAGE

[Turn over]



09.2

Sodium hydroxide solution was added gradually from a burette to 25 cm³ of 0.080 mol dm⁻³ propanoic acid at 25 °C

The pH was measured and recorded at regular intervals.

The results are shown in FIGURE 4, on page 76.

Use FIGURE 4 to determine the value of K_a for propanoic acid at 25 °C

Show your working. [3 marks]

K_a _____ mol dm^{-3}

[Turn over]



BLANK PAGE



09.3

Suggest which indicator is the most appropriate for the reaction in Question 09.2

Tick (✓) ONE box. [1 mark]

| Tick (✓) one box | Indicator | pH range |
|-----------------------------|-------------------------|--------------------|
| | methyl orange | 3.1 – 4.4 |
| | bromothymol blue | 6.0 – 7.6 |
| | cresolphthalein | 8.2 – 9.8 |
| | indigo carmine | 11.6 – 13.0 |

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 9 | . | 4 |
|---|---|---|---|

A student prepared a buffer solution by adding 0.0136 mol of a salt KX to 100 cm³ of a 0.500 mol dm⁻³ solution of a weak acid HX and mixing thoroughly.

The student then added 3.00×10^{-4} mol of potassium hydroxide to the buffer solution.

Calculate the pH of the buffer solution after adding the potassium hydroxide.

For the weak acid HX at 25 °C the value of the acid dissociation constant, $K_a = 1.41 \times 10^{-5}$ mol dm⁻³.

Give your answer to two decimal places.

[6 marks]



pH _____

[Turn over]



| | | | |
|---|---|---|---|
| 0 | 9 | . | 5 |
|---|---|---|---|

A buffer solution has a constant pH even when diluted.

Use a mathematical expression to explain this. [1 mark]

END OF QUESTIONS

| |
|----|
| |
| 15 |



BLANK PAGE



BLANK PAGE

| For Examiner's Use | |
|--------------------|------|
| Question | Mark |
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| 8 | |
| 9 | |
| TOTAL | |

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.

IB/M/JW/Jun19/7405/1/E4