



Surname _____

Other Names _____

Centre Number _____

Candidate Number _____

Candidate Signature _____

A-level

CHEMISTRY

Paper 2 Organic and Physical Chemistry

7405/2

Tuesday 11 June 2019 Afternoon

Time allowed: 2 hours

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.

[Turn over]



For this paper you must have:

- **the Periodic Table/Data Sheet, provided as an insert (enclosed)**
- **a ruler with millimetre measurements**
- **a scientific calculator, which you are expected to use where appropriate.**

INSTRUCTIONS

- **Use black ink or black ball-point pen.**
- **Answer ALL questions.**
- **You must answer the questions in the spaces provided. Do NOT write on blank pages.**
- **All working must be shown.**
- **Do all rough work in this book. Cross through any work you do not want to be marked.**



INFORMATION

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 105.**

DO NOT TURN OVER UNTIL TOLD TO DO SO



BLANK PAGE



Answer ALL questions in the spaces provided.

0 1

This question is about amines.

0 1 . 1

Give an equation for the preparation of 1,6-diaminohexane by the reaction of 1,6-dibromohexane with an excess of ammonia. [2 marks]

[Turn over]




0	1	.	2
---	---	---	---

On the opposite page, complete the mechanism for the reaction of ammonia with 6-bromohexylamine to form 1,6-diaminohexane.

Suggest the structure of a cyclic secondary amine that can be formed as a by-product in this reaction. [4 marks]



Mechanism



Cyclic secondary amine

[Turn over]



0 1 . 3

1,6-Diaminohexane can also be formed in a two-stage synthesis starting from 1,4-dibromobutane.

**Suggest the reagent and a condition for each stage in this alternative synthesis.
[3 marks]**

Stage 1 reagent and condition

Stage 2 reagent and condition



0	1	.	5
---	---	---	---

Justify the statement that there are no chiral centres in 3-aminopentane.
[1 mark]

12



BLANK PAGE

[Turn over]




0	2
---	---

A student prepared cyclohexene by heating cyclohexanol with concentrated phosphoric acid. The cyclohexene produced was distilled off from the reaction mixture.

0	2	.	1
---	---	---	---

Complete the diagram, on the opposite page, of the apparatus used to distil the cyclohexene from the reaction mixture at 83 °C. [2 marks]





[Turn over]



0 2 . 2

The distillate was shaken with saturated sodium chloride solution. The cyclohexene was separated from the aqueous solution using a separating funnel.

State why cyclohexene can be separated from the aqueous solution using the separating funnel. [1 mark]



02.3

The cyclohexene separated in Question 02.2 was obtained as a cloudy liquid. The student dried this cyclohexene by adding a few lumps of anhydrous calcium chloride and allowing the mixture to stand.

Give ONE observation that the student made to confirm that the cyclohexene was dry. [1 mark]

[Turn over]



0 2 . 4

In this preparation, the student added an excess of concentrated phosphoric acid to 14.4 g of cyclohexanol ($M_r = 100.0$).

The student obtained 4.15 cm³ of cyclohexene ($M_r = 82.0$).

Density of cyclohexene = 0.810 g cm⁻³

On the opposite page, calculate the percentage yield of cyclohexene obtained.

Give your answer to the appropriate number of significant figures. [5 marks]



% yield _____



[Turn over]



0	2	.	5
---	---	---	---

Cyclohexene reacts with bromine.

Complete the mechanism for this reaction. [3 marks]



12



BLANK PAGE

[Turn over]



0	3
---	---

The outer layers of some golf balls are made from a polymer called polyisoprene.

The isoprene monomer is a non-cyclic branched hydrocarbon that contains 88.2 % carbon by mass.

The empirical formula of isoprene is the same as its molecular formula.

0	3	.	1
---	---	---	---

Deduce the molecular formula of isoprene and, on the opposite page, suggest a possible structure. [4 marks]

Molecular formula _____




Structure

[Turn over]



03.2

The insides of some golf balls are made from a mixture of three other polymers. The repeating unit for one of these polymers is shown.



On the opposite page, draw the skeletal formula of the monomer used to make this polymer.

Give the IUPAC name of the monomer.
[2 marks]



Skeletal formula of monomer


IUPAC name _____

[Turn over]



0	3	.	3
---	---	---	---

A second polymer in the mixture has a repeating unit with the structure shown.



The third polymer in the mixture is a stereoisomer of this polymer.

Draw the structure of the repeating unit of the third polymer.

Give a reason why this type of stereoisomerism arises. [2 marks]

Repeating unit

Reason _____

[Turn over]



0	3	.	4
---	---	---	---

Golf balls recovered from lakes and ponds can be used again even after being in water for several years.

Explain why these golf balls do not biodegrade. [1 mark]

9



BLANK PAGE

[Turn over]



0 4

Substances P and Q react in solution at a constant temperature.

The initial rate of reaction was studied in three experiments by measuring the change in concentration of P over the first five seconds of the reaction.

The data obtained are shown in TABLE 1, on the opposite page.

28



TABLE 1

Experiment	Time after mixing / s	Concentration / mol dm⁻³	
		P	Q
1	0	1.00×10^{-2}	1.25×10^{-2}
	5.0	0.92×10^{-2}	not measured
2	0	2.00×10^{-2}	1.25×10^{-2}
	5.0	1.84×10^{-2}	not measured
3	0	0.50×10^{-2}	2.50×10^{-2}
	5.0	0.34×10^{-2}	not measured

[Turn over]

BLANK PAGE



04.1

Complete TABLE 2 to show the initial rate of reaction of P in each experiment. [1 mark]

TABLE 2

Experiment	Initial rate / mol dm ⁻³ s ⁻¹
1	1.6×10^{-4}
2	
3	

[Turn over]



04.2

Determine the order of reaction with respect to P and the order of reaction with respect to Q. [2 marks]

Order with respect to P _____

Order with respect to Q _____



BLANK PAGE

[Turn over]



04.3

A reaction between substances R and S was second order with respect to R and second order with respect to S. At a given temperature, the initial rate of reaction was $1.20 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$ when the initial concentration of R was $1.00 \times 10^{-2} \text{ mol dm}^{-3}$ and the initial concentration of S was $2.45 \times 10^{-2} \text{ mol dm}^{-3}$

Calculate a value for the rate constant, k , for the reaction at this temperature. Give the units for k [3 marks]

k _____

Units _____

[Turn over]

6



0	5
---	---

The rate constant, k , for a reaction varies with temperature as shown by the equation

$$k = Ae^{-E_a/RT}$$

For this reaction, at 25 °C, $k = 3.46 \times 10^{-8} \text{ s}^{-1}$

The activation energy $E_a = 96.2 \text{ kJ mol}^{-1}$

The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

Calculate a value for the Arrhenius constant, A , for this reaction.
Give the units for A . [4 marks]

A _____

Units _____

[Turn over]

4



0	6
---	---

This question is about isomers.

0	6	.	1
---	---	---	---

Give a reagent and observations for a test-tube reaction to distinguish between 2-methylbutan-1-ol and 2-methylbutan-2-ol. [3 marks]

Reagent _____

Observation with 2-methylbutan-1-ol



Observation with 2-methylbutan-2-ol

[Turn over]

0	6	.	2
---	---	---	---

Compounds A and B both have the molecular formula $C_4H_8Br_2$

A has a singlet, a triplet and a quartet in its 1H NMR spectrum.

B has only two singlets in its 1H NMR spectrum.

Draw a structure for each of A and B.
[2 marks]

A

B



0	6	.	3
---	---	---	---

Compounds C and D both have the molecular formula $C_6H_3Br_3$

C has two peaks in its ^{13}C NMR spectrum.

D has four peaks in its ^{13}C NMR spectrum.

Draw a structure for each of C and D.
[2 marks]

C


D

[Turn over]




06.4

Compounds E, F, and G are isomers.




E



F





G

FIGURE 1, on pages 45 to 47, shows the infrared spectra of these isomers, but not necessarily in the same order.

Label each spectrum with the correct letter E, F or G in the box. [1 mark]

[Turn over]




BLANK PAGE



FIGURE 1

Transmittance
/ %




45

[Turn over]



4 5


FIGURE 1 continued



Transmittance

/ %

100



50

0

4000

3000

2000

1500


1000

500

Wavenumber / cm^{-1}



FIGURE 1 continued



Transmittance

/ %

100

50

0

4000

3000

2000

1500

1000

500

Wavenumber / cm⁻¹

[Turn over]




4 7




0	7
---	---

Isomers X and Y have the molecular formula C_5H_8O

Isomer X



Isomer Y



0	7	.	1
---	---	---	---

Give the IUPAC name for isomer X.
[1 mark]



07.2

Explain how and why isomers X and Y can be distinguished by comparing EACH of their


- **boiling points**
- **^{13}C NMR spectra**
- **infrared spectra.**

Use data from Tables A and C in the Data Booklet in your answer. [6 marks]

08

Paracetamol is a medicine commonly used to relieve mild pain.

Traditionally, paracetamol has been made industrially in a three-step synthesis from phenol.



08.1

Name the mechanism of the reaction in Step 1. [1 mark]

[Turn over]




BLANK PAGE



08.2

Complete the equation for the reaction in Step 2.
[1 mark]



[Turn over]




08.3

In theory, either ethanoyl chloride or ethanoic anhydride could be used in Step 3.

Complete the mechanism for the reaction of 4-aminophenol with ethanoyl chloride.

RNH₂ is used to represent 4-aminophenol in this mechanism.
[2 marks]



0	8	.	4
---	---	---	---

In practice, ethanoic anhydride is used in the industrial synthesis rather than ethanoyl chloride.

**Give ONE reason why ethanoyl chloride is NOT used in the industrial synthesis.
[1 mark]**

[Turn over]



BLANK PAGE



0	8	.	5
---	---	---	---

In Step 3 other aromatic products are formed as well as paracetamol.

Draw the structure of ONE of these other aromatic products. [1 mark]


[Turn over]



08.6



Chemists have recently developed a two-step process to produce paracetamol from phenol.

In the first step, phenol is oxidised to hydroquinone.



In the second step, hydroquinone reacts with ammonium ethanoate to form paracetamol.

On page 61, complete the equation for this second step.
[1 mark]



hydroquinone

paracetamol

[Turn over]



BLANK PAGE



08.6

Calculate the mass, in kg, of hydroquinone ($M_r = 110.0$) needed to produce 250 kg of paracetamol. [3 marks]

63

Mass _____ kg

[Turn over]

10




0	9
---	---

This question is about thin-layer chromatography (TLC).

- **A protein was hydrolysed to form a mixture of amino acids.**
- **A spot of this mixture was added to a TLC plate and the plate placed vertically in a small volume of solvent 1.**
- **When the solvent front reached nearly to the top of the plate, the plate was removed and allowed to dry.**
- **The plate was turned anticlockwise through 90° and placed vertically in a small volume of solvent 2.**
- **When the solvent front reached nearly to the top of the plate, the plate was again removed and allowed to dry.**
- **FIGURE 2, on the opposite page, shows the final TLC plate.**



FIGURE 2

**KEY**

- Spot seen after use of solvent 1
- Spot seen after use of solvent 2

[Turn over]



BLANK PAGE



09.1


Suggest a suitable reagent for the hydrolysis of a protein. [1 mark]

09.2

Suggest how the positions of the amino acids on the TLC plate were located. [1 mark]



Repeat of FIGURE 2

**KEY**

- Spot seen after use of solvent 1
- Spot seen after use of solvent 2

0 9 . 3

Deduce the minimum number of amino acids present in the original mixture.

[1 mark]

0 9 . 4

Suggest why it was necessary to use two different solvents. [1 mark]

[Turn over]

4



1	0
---	---

Some compounds with different molecular formulas have the same relative molecular mass to the nearest whole number.

1	0	.	1
---	---	---	---

A dicarboxylic acid has a relative molecular mass of 118, to the nearest whole number.

On the opposite page, deduce the molecular formula of the acid.

[3 marks]

Molecular formula _____

[Turn over]



10.2

A student dissolved some of the dicarboxylic acid from Question 10.1 in water and made up the solution to 250 cm³ in a volumetric flask.

In a titration, a 25.0 cm³ sample of the acid solution needed 21.60 cm³ of 0.109 mol dm⁻³ sodium hydroxide solution for neutralisation.

On the opposite page, calculate the mass, in g, of the dicarboxylic acid used. Give your answer to the appropriate number of significant figures. [4 marks]



Mass _____ g

[Turn over]



1	0	.	4
---	---	---	---

Draw the structure of a different diol also with molecular formula $C_6H_{14}O_2$ that has a 1H NMR spectrum that consists of two singlet peaks. [1 mark]

[Turn over]



BLANK PAGE



10.5

The dicarboxylic acid in question 10.1 and the isomers of $C_6H_{14}O_2$ in Questions 10.3 and 10.4 all have a relative molecular mass of 118

State why the dicarboxylic acid can be distinguished from the two diols by high resolution mass spectrometry using electrospray ionisation. [1 mark]

[Turn over]

10



1 1

This question is about esters including biodiesel.

1 1 . 1


An ester is formed by the reaction of an acid anhydride with $\text{CH}_3\text{CH}_2\text{OH}$

Complete the equation, on the opposite page. In your answer show clearly the structure of the ester.

**Give the IUPAC name of the ester.
[3 marks]**



Equation



Name of ester _____

[Turn over]



1 1 . 2

In a reaction to form biodiesel, one mole of a vegetable oil reacts with an excess of methanol to form two moles of an ester with molecular formula $C_{19}H_{34}O_2$ and one mole of an ester with molecular formula $C_{19}H_{36}O_2$

On page 81, draw the structure of the vegetable oil showing clearly the ester links.

You should represent the hydrocarbon chains in the form $C_x H_y$ where x and y are the actual numbers of carbon and hydrogen atoms. [2 marks]

80



[Turn over]



11.3

The compound $C_{19}H_{34}O_2$ is the methyl ester of *Z,Z*-octadeca-9,12-dienoic acid.


Part of the structure of the acid is shown.

On page 83, complete the skeletal formula to show the next part of the hydrocarbon chain to carbon atom number 14.

In your answer, show the *Z* stereochemistry around both $C=C$ double bonds. [2 marks]

82





11.4

83

Give an equation for the complete combustion of the ester $C_{19}H_{34}O_2$ [1 mark]

[Turn over]



1 1 . 5


Combustion of biodiesel produces greenhouse gases such as carbon dioxide that cause global warming. Part of the infrared spectrum of carbon dioxide is shown in FIGURE 3, on the opposite page.

State how the infrared spectrum of carbon dioxide in FIGURE 3 is NOT what you might predict from the data provided in TABLE A in the Data Booklet. [1 mark]



FIGURE 3

Infrared spectrum of carbon dioxide



85

[Turn over]



BLANK PAGE



1 1 . 6

**Explain how carbon dioxide causes global warming.
[2 marks]**

[Turn over]

11



1 2

FIGURE 4, on page 89, shows two complementary strands in part of a DNA double helix structure.

1 2 . 1

Draw all the hydrogen bonds between the complementary strands shown in FIGURE 4.

**Use dashed lines to show the hydrogen bonds.
You do NOT need to show lone pairs of electrons or partial charges. [2 marks]**




1 2 . 2

Draw a ring around each of the component parts that make up the cytosine nucleotide in the section of DNA shown in FIGURE 4. [2 marks]



FIGURE 4



68

[Turn over]



BLANK PAGE



1 2 . 3

State the meaning of the term complementary when it is used to refer to DNA strands. [1 mark]

5

[Turn over]



1	3
---	---

Aqueous NaBH_4 reduces aldehydes but does not reduce alkenes.

1	3	.	1
---	---	---	---

On the opposite page, show the first step of the mechanism of the reaction between NaBH_4 and 2-methylbutanal.

You should include two curly arrows.

Explain why NaBH_4 reduces 2-methylbutanal but has no reaction with 2-methylbut-1-ene. [5 marks]



First step of mechanism

Explanation

[Turn over]



1	3	.	2
---	---	---	---

A student attempted to reduce a sample of 2-methylbutanal but added insufficient NaBH_4

The student confirmed that the reduction was incomplete by using a chemical test.

Give the reagent and observation for the chemical test. [2 marks]

Reagent _____

Observation _____

END OF QUESTIONS

7



BLANK PAGE



BLANK PAGE

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
TOTAL	

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.

IB/M/CD/Jun19/7405/2/E2