



**Surname** \_\_\_\_\_

**Other Names** \_\_\_\_\_

**Centre Number** \_\_\_\_\_

**Candidate Number** \_\_\_\_\_

**Candidate Signature** \_\_\_\_\_

**A-level**

**PHYSICS**

**Paper 2**

**7408/2**

**Friday 24 May 2019      Morning**

**Time allowed: 2 hours**

**At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.**

**[Turn over]**



**For this paper you must have:**

- **a pencil and a ruler**
- **a scientific calculator**
- **a Data and Formulae Booklet.**

## **INSTRUCTIONS**

- **Use black ink or black ball-point pen.**
- **Answer ALL questions.**
- **You must answer the questions in the spaces provided. Do not write on blank pages.**
- **If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).**
- **Do all rough work in this book. Cross through any work you do not want to be marked.**
- **Show all your working.**

## **INFORMATION**

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 85.**
- **You are expected to use a scientific calculator where appropriate.**
- **A Data and Formulae Booklet is provided as a loose insert.**

**DO NOT TURN OVER UNTIL TOLD TO DO SO**

## SECTION A

Answer ALL questions in this section.

01


**FIGURE 1** shows a perfectly insulated cylinder containing 0.050 kg of liquid nitrogen at a temperature of 70 K.

A heater transfers energy at a constant rate of 12 W to the nitrogen.

A piston maintains the pressure at  $1.0 \times 10^5$  Pa during the heating process.

**FIGURE 1**

**NOT TO SCALE**



**[Turn over]**



0	1	.	1
---	---	---	---

**The nitrogen is heated from 70 K and is completely turned into a gas after 890 s.**

**Calculate the specific heat capacity of liquid nitrogen.**

**Give an appropriate unit for your answer.**

**specific latent heat of vaporisation of nitrogen =  $2.0 \times 10^5 \text{ J kg}^{-1}$**

**boiling point of nitrogen = 77 K**

**[5 marks]**

**specific heat capacity = \_\_\_\_\_**

**Unit = \_\_\_\_\_**

**[Turn over]**



0	1	.	2
---	---	---	---

The work done by the nitrogen in the cylinder when expanding due to a change of state is X.

The energy required to change the state of the nitrogen from a liquid to a gas is Y.

Deduce which is greater, X or Y.

density of liquid nitrogen at its boiling temperature =  $810 \text{ kg m}^{-3}$

density of nitrogen gas at its boiling temperature =  $3.8 \text{ kg m}^{-3}$

[4 marks]



[Turn over]



0	2	.	1
---	---	---	---

**State what is meant by the internal energy of a gas. [2 marks]**

---

---

---

---

---

0	2	.	2
---	---	---	---

**Absolute zero of temperature can be interpreted in terms of the ideal gas laws or the kinetic energy of particles in an ideal gas.**

**Describe these two interpretations of absolute zero of temperature. [2 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



0	2	.	3
---	---	---	---

**A mixture of argon atoms and helium atoms is in a cylinder enclosed with a piston. The mixture is at a temperature of 310 K.**

**Calculate the root mean square speed ( $c_{\text{rms}}$ ) of the argon atoms in the mixture.**

**molar mass of argon =  $4.0 \times 10^{-2} \text{ kg mol}^{-1}$**

**[3 marks]**

13

$$c_{\text{rms}} = \underline{\hspace{10cm}} \text{ m s}^{-1}$$

**[Turn over]**



0	2	.	4
---	---	---	---

**Compare the mean kinetic energy of the argon atoms and the helium atoms in the mixture. [1 mark]**

---

---

---

---

---

0	2	.	5
---	---	---	---

**Explain, in terms of the kinetic theory model, why a pressure is exerted by the gas on the piston. [3 marks]**

---

---

---

---

15

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**[Turn over]**



0	2	.	6
---	---	---	---

**The mixture of gases in the cylinder stays the same.**

**Explain, using the kinetic theory model, TWO changes that can be made independently to reduce the pressure exerted by the gas. [3 marks]**

This image shows a blank sheet of white paper with ten horizontal black lines. The lines are evenly spaced and run across the width of the page, providing a template for writing or drawing.



---

---

---

---

---

---

---

---

---

---

[Turn over]



**BLANK PAGE**



0	3	.	1
---	---	---	---

**Define gravitational potential at a point.**  
**[1 mark]**

---

---

---

---

---


---

---

**[Turn over]**

FIGURE 2 shows the positions of equipotential surfaces at different distances from the centre of the Moon.

FIGURE 2 NOT TO SCALE



**Explain how the equipotential surfaces in FIGURE 2 show that the gravitational field is NOT uniform. [1 mark]**

---

---

---

---

---

---

---

---

**[Turn over]**



0	3	.	3
---	---	---	---


**Calculate, using FIGURE 2 on page 20, the escape velocity at the surface of the Moon.**

**radius of Moon =  $1.74 \times 10^6$  m**

**[4 marks]**

escape velocity = \_\_\_\_\_  $\text{m s}^{-1}$

[Turn over]



0	4
---	---

**FIGURE 3 shows an arrangement used to investigate the repulsive forces between two identical charged conducting spheres.**

**The spheres are suspended by non-conducting thread.**

**Each sphere has a mass of  $3.2 \times 10^{-3}$  kg and a radius of 20 mm.**

**The distance  $d$  is 40 mm.**


**The capacitance of a sphere of radius  $r$  is  $4\pi\epsilon_0 r$ .**

**Each sphere is charged by connecting it briefly to the positive terminal of a high-voltage supply, the other terminal of which is at 0 V.**

**After this has been done the charge on each sphere is 52 nC.**



FIGURE 3

NOT TO  
SCALE

[Turn over]



0	4	.	1
---	---	---	---

**Calculate the potential of one of the spheres. [3 marks]**

**potential = \_\_\_\_\_ V**

0	4	.	2
---	---	---	---

**The charged spheres in FIGURE 3, on page 25, are at equilibrium.**

**Draw labelled arrows on FIGURE 3 to show the forces on sphere B. [2 marks]**

0	4	.	3
---	---	---	---

**Suggest a solution to ONE problem involved in the measurement of  $d$  in FIGURE 3 on page 25. [2 marks]**

---

---

---

---

---

---

---

---

---

---

**[Turn over]**

0	4	.	4
---	---	---	---

**Show that the magnitude of the electrostatic force on each sphere is about  $4 \times 10^{-3}$  N. [3 marks]**

0	4	.	5
---	---	---	---

**A student measures the angle  $\theta$  when the apparatus in FIGURE 3, on page 25, is at equilibrium. The student records  $\theta$  as  $7^\circ$ .**

**Discuss whether this measurement is consistent with the other data in this investigation. [2 marks]**

---

---

---

---

---

---

---

**[Turn over]**



0	4	.	6
---	---	---	---

**The student says that the gravitational force between the two spheres has no **SIGNIFICANT** effect on the angle at which the spheres are in equilibrium.**

**Deduce with a calculation whether this statement is valid. [2 marks]**

---

---

---

[Turn over]




**A square coil of wire is rotating at a constant angular speed about a horizontal axis.**

**FIGURE 4 shows the coil at one instant when the normal to the plane of the coil is at  $30^\circ$  to a magnetic field.**

**The area of the coil is  $5.0 \times 10^{-4} \text{ m}^2$  and the flux density of the uniform magnetic field is  $2.5 \times 10^{-2} \text{ T}$ .**



**FIGURE 4**



**[Turn over]**

0	5	.	1
---	---	---	---

The maximum flux linkage of the coil during its rotation is  $1.5 \times 10^{-3}$  Wb turns.

Calculate the number of turns in the coil.  
[2 marks]

number of turns = \_\_\_\_\_



0	5	.	2
---	---	---	---

**Calculate the flux linkage of the coil at the instant shown in FIGURE 4 on page 33. [1 mark]**

**flux linkage = \_\_\_\_\_ Wb turns**

**[Turn over]**



05.3

The coil forms part of an electrical generator. FIGURE 5, on the opposite page, shows the emf generated by the coil.


Calculate the peak value of the emf generated. [2 marks]

36

emf = \_\_\_\_\_ V



FIGURE 5




05.4

**Sketch on FIGURE 6 the variation with time of flux linkage for the same time interval as FIGURE 5 on page 37.**  
**[1 mark]**

# FIGURE 6

flux linkage



39



3 9

[Turn over]

6

**BLANK PAGE**





0	6
---	---

**A thermal nuclear reactor uses a moderator to lower the kinetic energy of fast-moving neutrons.**

0	6	.	1
---	---	---	---

**Explain why the kinetic energy of neutrons must be reduced in a thermal nuclear reactor. [1 mark]**

---

---

---

---

---

---

---

**[Turn over]**

0	6	.	2
---	---	---	---

**As a result of a collision with an atom of a particular moderator, a neutron loses 63% of its kinetic energy.**

**A neutron has an initial kinetic energy of 2.0 MeV.**

**Calculate the kinetic energy of the neutron after five collisions. [2 marks]**

**kinetic energy = \_\_\_\_\_ eV**

**[Turn over]**



0	6	.	3
---	---	---	---

**The kinetic energy of a neutron in a thermal nuclear reactor is reduced from about 2 MeV to about 1 eV.**

**Explain why the number of collisions needed to do this depends on the nucleon number of the moderator atoms.  
[2 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

[Turn over]

0	6	.	4
---	---	---	---

One fission process which can occur in a thermal nuclear reactor is represented by the equation



Calculate in MeV the energy released in this fission process.

$$\text{mass of } {}_{92}^{235}\text{U} = 235.044 \text{ u}$$

$$\text{mass of } {}_{54}^{142}\text{Xe} = 141.930 \text{ u}$$

$$\text{mass of } {}_{38}^{90}\text{Sr} = 89.908 \text{ u}$$

$$\text{mass of } {}_0^1\text{n} = 1.0087 \text{ u}$$

[3 marks]



**energy released = \_\_\_\_\_ MeV**

**[Turn over]**



0	6	.	5
---	---	---	---

**Many magazine and newspaper articles focus on the risks of using nuclear power.**

**State THREE BENEFITS of using nuclear power. [3 marks]**

**1** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**2** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



3

---

---

---

---

---

[Turn over]

11

**SECTION B**

**Each of Questions 07 to 31 is followed by four responses, A, B, C and D.**

**For each question select the best response.**


**Only ONE answer per question is allowed.**

**For each question completely fill in the circle alongside the appropriate answer.**


**CORRECT METHOD** 

**WRONG METHODS**    

**If you want to change your answer you must cross out your original answer as shown.**



**If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.**



**You may do your working in the blank space around each question but this will not be marked.**

**Do NOT use additional sheets for this working.**

**[Turn over]**

0	7
---	---

## Brownian motion

[1 mark]

☐

**A makes it possible to see the motion of air molecules.**

☐

**B is caused by the collisions of smoke particles.**

☐

**C is caused by collisions between air molecules and smoke particles.**

☐

**D occurs because air is a mixture of gases and the molecules have different masses.**

08

Which row shows two scalar quantities?  
[1 mark]

☐

**A**

**gravitational  
potential**

**gravitational  
field strength**

☐

**B**

**mass**

**gravitational  
potential**

☐

**C**

**gravitational  
field strength**

**weight**

☐

**D**

**weight**

**gravitational  
potential**

[Turn over]

0	9
---	---

**What is the angular speed of a satellite in a geostationary orbit around the Earth?**  
**[1 mark]**

☐

**A**  $1.2 \times 10^{-5} \text{ rad s}^{-1}$

☐

**B**  $7.3 \times 10^{-5} \text{ rad s}^{-1}$

☐

**C**  $4.2 \times 10^{-3} \text{ rad s}^{-1}$

☐

**D**  $2.6 \times 10^{-1} \text{ rad s}^{-1}$

1	0
---	---

A planet of mass  $M$  and radius  $R$  rotates so quickly that material at its equator only just remains on its surface.

What is the period of rotation of the planet? [1 mark]

☐ **A**  $2\pi\sqrt{\frac{R}{GM}}$

☐ **B**  $2\pi\sqrt{\frac{GM}{R}}$

☐ **C**  $2\pi\sqrt{\frac{R^3}{GM}}$

☐ **D**  $2\pi\sqrt{\frac{GM}{R^3}}$

[Turn over]



1	1
---	---

**Satellites N and F have the same mass and are in circular orbits about the same planet. The orbital radius of F is greater than that of N.**

**Which is greater for F than for N?  
[1 mark]**

☐

**A gravitational force on the satellite**

☐

**B angular speed**

☐

**C kinetic energy**

☐

**D orbital period**



1	2
---	---

**An object moves freely at  $90^\circ$  to the direction of a gravitational field.**

**The acceleration of the object is**

**[1 mark]**

☐

**A zero.**

☐

**B opposite to the direction of the gravitational field.**

☐

**C in the direction of the gravitational field.**

☐

**D at  $90^\circ$  to the direction of the gravitational field.**


**[Turn over]**

1	3
---	---


When an electron is moving at a speed  $v$  perpendicular to a uniform magnetic field of flux density  $B$ , it follows a path of radius  $R$ .

A second electron moves at a speed  $\frac{v}{2}$  perpendicular to a uniform magnetic field of flux density  $4B$ .


**What is the radius of the path of the second electron? [1 mark]**




**A**  $\frac{R}{8}$



**B**  $\frac{R}{4}$



**C**  $2R$




**D**  $8R$

**[Turn over]**



1	4
---	---

A small object of mass  $m$  has a charge  $Q$ . The object remains stationary in an evacuated space between two horizontal plates. The plates are separated by a distance  $d$  and the potential difference between the plates is  $V$ .



What is  $V$ ? [1 mark]

☐

**A**  $\frac{m Q g}{d}$

☐

**B**  $\frac{m d g}{Q}$

☐

**C**  $\frac{m Q}{d}$

☐

**D**  $\frac{m d}{Q}$

[Turn over]



1	5
---	---

**1.5 mJ of work is done when a charge of  $30\ \mu\text{C}$  is moved between two points, M and N, in an electric field.**

**What is the potential difference between M and N? [1 mark]**

☐

**A    20 mV**

☐

**B    20 V**

☐

**C    45 V**

☐

**D    50 V**

1	6
---	---

**An electric field acts into the plane of the paper. An electron enters the field at  $90^\circ$  to the field lines.**

**The force on the electron is**

**[1 mark]**

☐

**A zero.**

☐

**B along the direction of the field.**

☐

**C at  $90^\circ$  to the field.**

☐

**D opposite to the direction of the field.**

**[Turn over]**



17

The ionisation potential for the atoms of a gas is  $V$ . Electrons of mass  $m$  and charge  $e$  travelling at a speed  $v$  can just cause ionisation of atoms in the gas.

What is  $v$ ? [1 mark]

☐

**A**  $\frac{eV}{2m}$

☐

**B**  $\frac{2eV}{m}$

☐

**C**  $\sqrt{\frac{eV}{2m}}$

☐

**D**  $\sqrt{\frac{2eV}{m}}$



**BLANK PAGE**

**[Turn over]**



1	8
---	---

**When a small radioactive source is placed in a cloud chamber, straight tracks about 4 cm long are observed. The same source is placed 10 cm from a Geiger tube and a count rate is detected. When a sheet of aluminium 5 mm thick is placed between the source and the Geiger tube the count rate falls to the background count rate.**

**Which types of radiation are emitted by the source? [1 mark]**

☐

**A**  $\alpha$ ,  $\beta$  and  $\gamma$

☐

**B**  $\beta$  and  $\gamma$

☐

**C**  $\alpha$  and  $\gamma$

☐

**D**  $\alpha$  and  $\beta$

**[Turn over]**

19

A parallel-plate capacitor is made by inserting a sheet of dielectric material between two plates. Both plates are in contact with the sheet.

Which relative permittivity and sheet thickness give the greatest capacitance?  
[1 mark]

	Relative permittivity	Thickness / mm
<input type="radio"/>	A 2	0.40
<input type="radio"/>	B 3	0.90
<input type="radio"/>	C 4	1.0
<input type="radio"/>	D 6	1.6

2	0
---	---

**A  $1.0 \mu\text{F}$  capacitor is charged for 20 s using a constant current of  $10 \mu\text{A}$ .**

**What is the energy transferred to the capacitor? [1 mark]**

☐

**A**  $5.0 \times 10^{-3} \text{ J}$

☐

**B**  $1.0 \times 10^{-2} \text{ J}$

☐

**C**  $2.0 \times 10^{-2} \text{ J}$

☐

**D**  $4.0 \times 10^{-2} \text{ J}$

**[Turn over]**

2	1
---	---

**A  $1.0\ \mu\text{F}$  capacitor initially stores  $15\ \mu\text{C}$  of charge. It then discharges through a  $25\ \Omega$  resistor.**

**What is the maximum current during the discharge of the capacitor? [1 mark]**

☐

**A     $0.60\ \text{mA}$**

☐

**B     $1.2\ \text{mA}$**

☐

**C     $0.60\ \text{A}$**

☐

**D     $1.2\ \text{A}$**

22

The initial potential difference across a capacitor is  $V_0$ . The capacitor discharges through a circuit of time constant  $T$ . The base of natural logarithms is  $e$ .

What is the potential difference across the capacitor after time  $T$ ? [1 mark]

☐

**A**  $\frac{V_0}{2}$

☐

**B**  $\frac{V_0}{e}$

☐

**C**  $V_0 e$

☐


**D**  $V_0 \ln 2$

[Turn over]



2	3
---	---

The plane of coil PQRS is parallel to a uniform magnetic field.





**When a current  $I$  is in the coil**

**[1 mark]**

☐

**A there are no magnetic forces acting on SP and QR.**

☐

**B there are no magnetic forces acting on PQ and RS.**

☐

**C an attractive magnetic force acts between SP and QR.**

☐

**D an attractive magnetic force acts between PQ and RS.**

**[Turn over]**

2	4
---	---

**A horizontal wire of length 0.50 m and weight 1.0 N is placed in a uniform horizontal magnetic field of flux density 1.5 T directed at  $90^\circ$  to the wire.**

**What is the current that just supports the wire? [1 mark]**

☐

**A    0.33 A**

☐

**B    0.75 A**

☐

**C    1.3 A**

☐

**D    3.0 A**

2	5
---	---

**Which is NOT an assumption about gas particles in the kinetic theory model for a gas? [1 mark]**

☐

**A They collide elastically with the container walls.**

☐

**B They have negligible size compared to the distance between the container walls.**

☐

**C They travel between the container walls in negligibly short times.**

☐

**D They collide with the container walls in negligibly short times.**


**[Turn over]**



2	6
---	---

**A coil P is connected to a cell and a switch.**

**A second closed coil Q is parallel to P and is arranged on the same axis.**



**When the switch is closed, coil Q experiences a force.**


Which row describes the force on Q?  
[1 mark]

		FORCE	DIRECTION OF FORCE
<input type="radio"/>	A	increases to constant value	to left
<input type="radio"/>	B	increases to constant value	to right
<input type="radio"/>	C	increases then decreases	to left
<input type="radio"/>	D	increases then decreases	to right

[Turn over]

2	7
---	---

**Three identical magnets P, Q and R are released simultaneously from rest and fall to the ground from the same height.**



**P falls directly to the ground.**

**Q falls through the centre of a thick horizontal conducting ring.**

**R falls through a similar ring that has a gap cut into it.**

**In which order do the magnets reach the ground? [1 mark]**

☐

**A P and R arrive together, followed by Q.**

☐

**B P and Q arrive together, followed by R.**

☐

**C P arrives first, followed by Q which is followed by R.**

☐

**D All three magnets arrive simultaneously.**

**[Turn over]**



2	8
---	---

A steady current  $I$  dissipates power  $P$  in a resistor of resistance  $R$ .

An alternating current through a resistor of resistance  $2R$  has a peak value of  $I$ .

What is the power dissipated in the second resistor? [1 mark]

☐

**A**  $\frac{P}{\sqrt{2}}$

☐

**B**  $P$

☐

**C**  $\sqrt{2} P$

☐

**D**  $2P$




**BLANK PAGE**

**[Turn over]**



The figure shows an oscilloscope trace of a sinusoidal ac voltage.



The time base setting is  $5\text{ ms cm}^{-1}$  and the Y-voltage gain is  $10\text{ V cm}^{-1}$ .

Which row describes the ac voltage? [1 mark]

☐

A

14

50

Frequency / Hz

☐

B

14

100

☐

C

7

50

☐

D

7


100



[Turn over]

3	0
---	---

**A deuterium nucleus and a tritium nucleus fuse together to form a helium nucleus and a particle X. The equation for this process is:**



**What is X? [1 mark]**

☐

**A electron**

☐

**B neutron**

☐

**C positron**

☐

**D proton**

31

What effect are the control rods intended to have on the average kinetic energy and number of fission neutrons in a thermal nuclear reactor? [1 mark]

		Average kinetic energy of fission neutrons	Number of fission neutrons
<input type="radio"/>	A	unchanged	unchanged
<input type="radio"/>	B	reduced	unchanged
<input type="radio"/>	C	unchanged	reduced
<input type="radio"/>	D	increased	reduced

END OF QUESTIONS



**Additional page, if required.**

**Write the question numbers in the left-hand margin.**


**Additional page, if required.**

**Write the question numbers in the left-hand margin.**

This image shows a blank sheet of white paper with horizontal ruling lines. A single vertical line runs down the left side, creating a narrow margin. There are 20 horizontal lines in total, evenly spaced across the page. The lines are thin and black.

**Additional page, if required.**

**Write the question numbers in the left-hand margin.**




**Additional page, if required.**

**Write the question numbers in the left-hand margin.**


BLANK PAGE

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7-31	
TOTAL	

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk) after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ

Copyright © 2019 AQA and its licensors. All rights reserved.

IB/M/JW/Jun19/7408/2/E2

