



**Surname** \_\_\_\_\_

**Other Names** \_\_\_\_\_

**Centre Number** \_\_\_\_\_

**Candidate Number** \_\_\_\_\_

**Candidate Signature** \_\_\_\_\_

**I declare this is my own work.**

**A-level**

**BIOLOGY**

**Paper 3**

**7402/3**

**Time allowed: 2 hours**

**At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.**

**[Turn over]**



**BLANK PAGE**



**For this paper you must have:**

- **a ruler with millimetre measurements**
- **a scientific calculator.**

## **INSTRUCTIONS**

- **Use black ink or black ball-point pen.**
- **Answer ALL questions in SECTION A.**
- **Answer ONE question from SECTION B.**
- **You must answer the questions in the spaces provided. Do not write on blank pages.**
- **If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).**
- **Show all your working.**
- **Do all rough work in this book. Cross through any work you do not want to be marked.**

## **INFORMATION**

- **The marks for the questions are shown in brackets.**
- **The maximum mark for this paper is 78.**

**DO NOT TURN OVER UNTIL TOLD TO DO SO**



## SECTION A

Answer ALL questions in this section.

You are advised to spend no more than 1 hour and 15 minutes on this section.

0	1
---	---

In one species of squirrel, ‘*Sciurus carolinensis*’, fur colour is controlled by one gene, with two codominant alleles.  $C^G$  represents the allele for grey fur colour, and  $C^B$  represents the allele for black fur colour.

TABLE 1 shows the three possible phenotypes.

TABLE 1

GENOTYPE	PHENOTYPE
$C^G C^G$	Grey fur
$C^G C^B$	Brown-black fur
$C^B C^B$	Black fur



0	1	.	1
---	---	---	---

In a population of 34 'S. carolinensis', 2 had black fur.

Use the Hardy–Weinberg equation to estimate how many squirrels in this population had brown-black fur. Show your working. [2 marks]

Answer \_\_\_\_\_

[Turn over]



0	1	.	2
---	---	---	---

The actual number of squirrels in this population that had brown-black fur was 16.

Use all of the information to calculate the ACTUAL frequency of the  $C^G$  allele.

Do NOT use the Hardy–Weinberg equation in your calculation.

Give your answer to 2 decimal places. [1 mark]

Answer \_\_\_\_\_



0	1	.	3
---	---	---	---

**‘S. carolinensis’ were first introduced to the UK from North America in the 1870s. They are now widely distributed across the UK.**

**‘S. carolinensis’ from both North America and the UK show exactly the same genotypic and phenotypic variation. An identical mutation causing black fur has also been found in several other species closely related to ‘S. carolinensis’.**

**Use this information to deduce which ONE of the following conclusions is most likely true.**

**Tick (✓) ONE box. [1 mark]**

☐

**A The mutation that caused black fur happened after ‘S. carolinensis’ was introduced to the UK from North America.**

☐

**B The mutation that caused black fur happened in a common ancestor of ‘S. carolinensis’ and other closely related species.**

☐

**C The mutation that caused black fur happened independently in ‘S. carolinensis’ and all other closely related species.**

☐

**D The phenotypic variation shown in ‘S. carolinensis’ and other closely related species is caused by genetic drift.**

**[Turn over]**



The mutation that caused the  $C^B$  allele was due to a 24 base-pair deletion from the  $C^G$  allele.

01.4

The protein coded for by the  $C^B$  allele is 306 amino acids long.

Calculate the percentage reduction in size of the protein coded for by the  $C^B$  allele compared with the protein coded for by the  $C^G$  allele.

Give your answer to 3 significant figures and show your working. [2 marks]

Answer \_\_\_\_\_




**BLANK PAGE**

**[Turn over]**



In '*S. carolinensis*', fur colour depends on the distribution and relative amounts of light pigments and dark pigments in the hairs of the fur. FIGURE 1 shows how the protein produced from the  $C^G$  allele can result in the production of a light pigment or a dark pigment.

**FIGURE 1**



The deletion mutation in the  $C^B$  allele results in the production of a receptor protein that does not have glutamic acid. The lack of glutamic acid in the receptor protein has the same effect as  $\alpha$ MSH leaving the receptor protein.

01.5

Use FIGURE 1 and this information to suggest why 'S. carolinensis' with the genotype  $C^B C^B$  have black fur rather than grey fur. [3 marks]

---

---

---

---

---

---

---

---

---

[Turn over]



---

---

---

---

---

---

<hr/>
9



**BLANK PAGE**

**[Turn over]**



0	2	.	1
---	---	---	---

**Describe how the human immunodeficiency virus (HIV) is replicated ONCE INSIDE helper T cells (T<sub>H</sub> cells).**

**[4 marks]**

[illegible]

---

---

---

---

---

**[Turn over]**



**HIV-1 is the most common type of HIV. HIV-1 binds to a receptor on T<sub>H</sub> cells called CCR5.**

**Current treatment for HIV-1 involves the use of daily antiretroviral therapy (ART) to stop the virus being replicated. Only 59% of HIV-positive individuals have access to ART.**

**Scientists have found that two HIV-1-positive patients (P and Q) have gone into remission (have no detectable HIV-1). This happened after a blood stem cell transplant (BSCT).**

- Patient P was given TWO BSCTs, and patient Q was given ONE BSCT.**
- All BSCTs came from a donor with T<sub>H</sub> cells WITHOUT the CCR5 receptor.**
- In addition, patient P had radiotherapy, and patient Q had chemotherapy. Both of these treatments are toxic.**
- BOTH patients (P and Q) stopped receiving ART 16 months after BSCT.**

**18 months after stopping ART, BOTH patients had NO HIV-1 RNA in their plasma, NO HIV-1 DNA in their T<sub>H</sub> cells and NO CCR5 on their T<sub>H</sub> cells.**



0	2	.	2
---	---	---	---

**Use the information given to evaluate the use of BSCT to treat HIV infections. [5 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

**[Turn over]**



---

---

---

---

---

---

---

---

---

<hr/>
9



**BLANK PAGE**

**[Turn over]**



0	3
---	---

**Scientists investigated movement in adult pine beetles. Adult beetles emerge from cracks in tree bark.**

**The scientists released a newly emerged adult beetle, G, from the centre of a sample area that had a single light source coming from one direction. They made a drawing of the beetle's path of walking. They repeated this with three more beetles, J, P and R.**

**FIGURE 2, on the opposite page, shows the scientists' results.**

0	3	.	1
---	---	---	---

**Name the type of behaviour shown by beetles G, J, P and R, and suggest ONE advantage to adult beetles of the type of behaviour shown. [2 marks]**

**Behaviour** \_\_\_\_\_

**Advantage** \_\_\_\_\_


---


---

---



FIGURE 2

**KEY**

-  **Position of light source**
- **Point from which beetle was released**

[Turn over]



**At higher temperatures and higher light intensities, adult pine beetles normally**


- **move more**
- **fly rather than walk.**

**When preparing to fly, these adult beetles walk slowly. The scientists investigated the movement of adult beetles at different temperatures, and in the light and the dark. They created a box that was half in the light and half in the dark. They released an adult beetle at the midpoint of the central dividing line between light and dark areas. They recorded the path of the beetle's movement and its location after 5 minutes. From this, they calculated the mean speed of movement. They repeated the experiment with many beetles and at several temperatures.**

**FIGURE 3, on the opposite page, shows the scientists' results.**



FIGURE 3



[Turn over]



**BLANK PAGE**



0	3	.	2
---	---	---	---

**After studying these experiments, a student concluded:**

- **there is a significant change in movement between 35 °C and 37.5 °C**
- **between 35 °C and 37.5 °C, more beetles move away from the light**
- **between 35 °C and 37.5 °C, more beetles have a slower walking speed.**

**Suggest reasons why these conclusions might NOT be valid. [3 marks]**

---

---

---

---

---

---

---

---

---

**[Turn over]**



---

---

---

---

---

<hr/>
5



**BLANK PAGE**

**[Turn over]**



0	4
---	---

Freshwater marshes have one of the highest rates of gross primary production (*GPP*) and net primary production (*NPP*) of all ecosystems.

Carbon use efficiency (*CUE*) is the ratio of *NPP:GPP*. Freshwater marshes have a high *CUE*.

0	4	.	1
---	---	---	---

Use your knowledge of *NPP* to explain why freshwater marshes have a high *CUE* AND the advantage of this.

Do NOT refer to abiotic factors in your answer.  
[2 marks]

Explanation \_\_\_\_\_

\_\_\_\_\_

Advantage \_\_\_\_\_

\_\_\_\_\_



0	4	.	2
---	---	---	---

**Freshwater marsh soils are normally waterlogged. This creates anaerobic conditions.**

**Use your knowledge of the nitrogen cycle to suggest why these soils contain relatively high concentrations of ammonium compounds and low concentrations of nitrite ions and nitrate ions. [2 marks]**

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



**A student investigated the growth rate of a freshwater marsh plant.**

**The growth rate ( $R$ ) of a plant can be determined using this equation.**

$$R = \frac{(\ln W_2 - \ln W_1)}{t}$$

**Where**

**$\ln$  = natural logarithm**

**$t$  = duration of the investigation in days**

**$W_1$  = plant biomass at the start of the investigation**

**$W_2$  = plant biomass at the end of the investigation**

**The student used the equation above; however, she substituted height for biomass. This was because she did not want to destroy the plants to measure their biomass.**



0	4	.	3
---	---	---	---

**State the assumption the student has made AND suggest why this assumption might NOT be valid.**  
**[2 marks]**

---

---

---

---

---

---

---

---

---

**[Turn over]**



0	4	.	4
---	---	---	---

At the end of the investigation, the student noted the freshwater marsh plant had grown 268 mm in height, and now measured 387 mm. She calculated the rate of growth ( $R$ ) to be  $0.097 \text{ mm m}^{-1} \text{ day}^{-1}$

Use this information and, **SUBSTITUTING HEIGHT FOR BIOMASS**, use the equation to calculate the duration of the student's investigation.

Give your answer to the nearest full day. Show your working. [2 marks]

\_\_\_\_\_ days

—
8



0	5	.	1
---	---	---	---

The action of endopeptidases and exopeptidases can increase the rate of protein digestion. Describe how.  
[2 marks]

---

---

---

---

---

---

---

---

---

---

[Turn over]



**05.2**

**As humans age, there is a decrease in body protein.**

**Give the name of ONE body protein that could have resulted in:**

**[2 marks]**

**reduced muscle power** \_\_\_\_\_

\_\_\_\_\_

**reduced immunity** \_\_\_\_\_

\_\_\_\_\_

**Scientists investigated the effect of two types of dietary protein on the ability of old men to produce body proteins.**

**TABLE 2, on the opposite page, shows information about the two types of dietary protein investigated.**



TABLE 2

Physiological factor	Name of dietary protein	
	Casein	Whey
Rate of absorption of dietary protein / $\text{mmol dm}^{-3}$ amino acids in blood plasma $\text{h}^{-1}$	3.05	4.33
Stimulation of protein synthesis	Higher rate	Lower rate
Breakdown of body proteins	No effect	Inhibitory effect


[Turn over]



**FIGURE 4** shows the percentage of protein absorbed that becomes body protein in old men following a meal of casein or whey.

**FIGURE 4**

Percentage  
of protein  
absorbed  
that becomes  
body protein



A statistical test confirmed that the difference between the results shown in **FIGURE 4** was significant.

05.3

**Suggest which type of dietary protein would be better for old men to eat to cause a NET gain of body proteins. Use the information provided to explain your answer. [3 marks]**

---

---



7



0	6
---	---

Plants transport sucrose from leaves to other tissues for growth and storage. SUT1 is a sucrose co-transporter protein.

Scientists investigated whether the cells of tobacco plant leaves used SUT1 to transport sucrose to other tissues.

0	6	.	1
---	---	---	---

The scientists used a radioactively labelled DNA probe to show that the cells of tobacco plant leaves contained the *SUT1* gene.

Describe how they would do this.

Do NOT include PCR in your answer. [4 marks]

---

---

---

---

---

---

---

---





**06.2**

To study the role of *SUT1* in tobacco plants, scientists reduced the expression of the *SUT1* gene.

When the *SUT1* gene is transcribed, the *SUT1* mRNA produced is called 'sense' *SUT1* mRNA. The scientists genetically modified plants by inserting an EXTRA gene so that this ALSO allowed the production of 'antisense' *SUT1* mRNA.

The scientists had two types of tobacco plants:

- type A – plants that were genetically modified
- type B – plants that were NOT genetically modified.

Suggest how the production of 'antisense' *SUT1* mRNA in type A plants would reduce the expression of the *SUT1* gene. [4 marks]

---

---

---

---

---

---

---





0	6	.	3
---	---	---	---

The scientists hypothesised that lower rates of sucrose transport from leaves would cause reduced growth.

To test this hypothesis, the scientists provided leaves of type A and type B plants with labelled carbon dioxide ( $^{14}\text{CO}_2$ ). To estimate sucrose transport out of leaves, they measured the percentage of  $^{14}\text{C}$  remaining in the leaves for 16 hours.

FIGURE 5, on the opposite page, shows their results.


Calculate the ratio of percentage of  $^{14}\text{C}$  remaining in leaves of type B to type A plants 16 hours after providing  $^{14}\text{CO}_2$  [1 mark]

Answer \_\_\_\_\_



FIGURE 5

Percentage of  
 $^{14}\text{C}$  remaining  
in the leaves



KEY


- Type A plants  
— Type B plants

[Turn over]



## REPEAT OF FIGURE 5

Percentage of  
 $^{14}\text{C}$  remaining  
in the leaves



KEY

- Type A plants  
— Type B plants



0	6	.	4
---	---	---	---

In type B plants, the percentage of  $^{14}\text{C}$  remaining in the leaves does not reach zero per cent, as shown in FIGURE 5.

Suggest TWO reasons why. [2 marks]

1 \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

2 \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

[Turn over]



The scientists measured physiological differences between type A plants and type B plants.

TABLE 3 shows the scientists' results as they presented them.

**TABLE 3**

Physiological factor	Type of tobacco plant	
	Type A	Type B
Rate of sucrose transport from leaf cells / $\mu\text{mol m}^{-2} \text{ s}^{-1}$	0.1	3.7
Leaf sucrose concentration / $\text{mmol m}^{-2}$	22	4
Ratio of shoot:root dry mass	6:1	2:1
Rate of photosynthesis / $\mu\text{mol glucose m}^{-2} \text{ s}^{-1}$	4	14

Sucrose is able to inhibit the production and activity of rubisco in leaves of a plant. Type A plants have decreased dry mass compared with type B plants.



0	6	.	5
---	---	---	---

**Use all the information to suggest AND explain how the physiological factors in TABLE 3 would contribute to the decreased dry mass observed in type A plants.  
[4 marks]**

[illegible]

**[Turn over]**



This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

15



**BLANK PAGE**

**[Turn over]**



**SECTION B**

**Answer ONE question.**

**You are advised to spend no more than 45 minutes on this section.**

0	7
---	---

**Write an essay on ONE of the topics below.**

**EITHER**

0	7	.	1
---	---	---	---

**The importance of complementary shapes of molecules in organisms [25 marks]**

**OR**

0	7	.	2
---	---	---	---

**The importance of ions in metabolic processes [25 marks]**

---

---





















[illegible]



























[illegible]





25



**Additional page, if required.**

**Write the question numbers in the left-hand margin.**

[illegible]

**Additional page, if required.**

**Write the question numbers in the left-hand margin.**

[illegible]

**BLANK PAGE**

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
<b>TOTAL</b>	

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk).

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2021 AQA and its licensors. All rights reserved.

**IB/M/CD/Jun21/7402/3/E2**