

Surname
Other Names
Centre Number
Candidate Number
Candidate Signature
I declare this is my own work.

AS

CHEMISTRY

Paper 1 Inorganic and Physical Chemistry

7404/1

Time allowed: 1 hour 30 minutes

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.



For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Answer ALL questions.
- You must answer the questions in the spaces provided. Do NOT write on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.



INFORMATION

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

ADVICE

You are advised to spend about 65 minutes on SECTION A and 25 minutes on SECTION B.

DO NOT TURN OVER UNTIL TOLD TO DO SO



SECTION A

Answer ALL questions in this section.

0 1

This question is about ionisation energies of Group 2 elements.

0 1 . 1

Explain why the first ionisation energy of the Group 2 elements decreases down the group. [2 marks]



01.2
Give an equation, including state symbols, to represent the process that occurs when the third ionisation energy of magnesium is measured. [1 mark]
0 1 . 3
Explain why the third ionisation energy of magnesium is much higher than the second ionisation energy of magnesium. [2 marks]
[Turn over]5



0	2


This question is about acid-base titrations.

Citric acid reacts with sodium hydroxide.

$$C_6H_8O_7(aq) + 3NaOH(aq) \longrightarrow$$
 $Na_3C_6H_5O_7(aq) + 3H_2O(l)$

A student makes a solution of citric acid by dissolving some solid citric acid in water.

Describe a method to add an accurately known mass of solid to a beaker to make a solution. [2 marks]





0	2		2
_	_	- 1	_

The student dissolves 0.834 g of citric acid in water and makes the solution up to 500 $\,\mathrm{cm}^3$

Calculate the concentration, in mol dm^{-3} , of citric acid in this solution. [3 marks]

Concentration	mol dm ⁻³



02.3

The student uses this method to complete a titration.

- Rinse a burette with distilled water.
- Fill the burette with sodium hydroxide solution.
- Use a measuring cylinder to transfer 25 cm³ of the citric acid solution into a conical flask.
- Add 5 cm³ of indicator.
- Slowly add the sodium hydroxide solution from the burette into the conical flask.
- Add the sodium hydroxide solution dropwise near the end point until the indicator just changes colour.
- Repeat the titration to get concordant results.

The method used by the student includes three practical steps that will lead to an inaccurate final result.

For each of these three steps

- identify the mistake
- explain why it is a mistake
- suggest how the mistake can be overcome.

[6 marks]











BLANK PAGE



02.4

TABLE 1 shows the student's burette readings after the mistakes in the practical procedure have been corrected.

TABLE 1

	ROUGH	RUN 1	RUN 2	RUN 3
FINAL READING / cm ³	23.65	22.95	46.05	26.30
START READING / cm ³	0.00	0.00	22.95	3.40
TITRE / cm ³	23.65			

Complete TABLE 1.

Use the data in TABLE 1 to calculate the mean titre. [2 marks]

Mean	titre	С	m ³



[Turn over]
Percentage uncertainty
Calculate the percentage uncertainty in the use of the burette in RUN 1. [1 mark]
0 2 . 5 The total uncertainty in the use of the burette is ±0.15 cm ³



BLANK PAGE



0 3

This question is about shapes of molecules.

Complete TABLE 2 by drawing the shapes of both the AsF_5 and KrF_2 molecules, showing all lone pairs of electrons that influence the shape.

Deduce the bond angle(s) in AsF₅ [3 marks]

TABLE 2

	AsF ₅	KrF ₂
DIAGRAM OF SHAPE		
BOND ANGLE(S)		

Turn over



3

0 4

This question is about intermolecular forces.

04.1

Complete the diagram to show how one molecule of ammonia can form a hydrogen bond with one molecule of ethanol.

Include all lone pairs of electrons and partial charges on atoms involved in the hydrogen bond. [3 marks]






TABLE 3 shows the electronegativity values of atoms of some elements.

TABLE 3

ATOM	Н	С	N	0	Br
ELECTRONEGATIVITY	2.1	2.5	3.0	3.5	2.8

0 4		2	
-----	--	---	--

Define the	term	electronegativity.	[1	mark]
------------	------	--------------------	----	-------

04.3

Deduce the TWO atoms from TABLE 3 that will form the most polar bond. [1 mark]



04.4
The C-Br bond is polar.
Explain why CBr ₄ is NOT a polar molecule. [2 marks]



04.5	
Suggest, in terms of the intermolecular forces for eacompound, why CBr ₄ has a higher boiling point than CHBr ₃ [3 marks]	
	_
[Turn over]	
	10



0 5

A sample of antimony is analysed in a time of flight (TOF) mass spectrometer and is found to contain two isotopes, ¹²¹Sb and ¹²³Sb

After electron impact ionisation, all of the ions are accelerated to the same kinetic energy (KE) and then travel through a flight tube that is 1.05 m long. A 121 Sb⁺ ion takes 5.93×10^{-4} s to travel through the flight tube.

The kinetic energy of an ion is given by the equation $KE = \frac{1}{2}mv^2$

KE = kinetic energy / J m = mass / kg v = speed / m s⁻¹

Calculate the mass, in kg, of one ¹²¹Sb⁺ ion.

Calculate the time taken for a ¹²³Sb⁺ ion to travel through the same flight tube.

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$ [5 marks]



Mass of one ¹²¹ Sb ⁺ ion	kg
--	----

Time taken by a ¹²³Sb⁺ ion _____ s

[Turn over]

5



0	6
•	

lodide ions can be oxidised to iodine using oxidising agents such as iodate(V) ions (IO_3^-) and concentrated sulfuric acid.

0 6		1
-----	--	---

State, in terms of electrons, the meaning of the term oxidising agent. [1 mark]



In acidic solution, IO_3^- ions oxidise iodide ions to iodine.

$$10_3^- + 51^- + 61^+ \rightarrow 31_2 + 31_20$$

Give a half-equation for the oxidation of iodide ions to iodine.

Deduce the half-equation to show the reduction process in this reaction. [2 marks]

Oxidation half-equation

Reduction half-equation



06.3
When iodide ions are oxidised using concentrated sulfuric acid, sulfur dioxide, a yellow solid and a foul-smelling gas are all formed.
Give an equation to show the reaction between iodide ions and concentrated sulfuric acid to form the yellow solid

Identify the foul-smelling gas. [2 marks]

Equation

Identity of foul-smelling gas

5



BLANK PAGE




0 7

This question is about gaseous equilibria.

FIGURE 1 shows the effect of pressure on the percentage yield of a reaction at equilibrium at two different temperatures.

FIGURE 1







07.1			
Explain how FIGURE 1 shows that the forward reaction in this equilibrium is exothermic. [2 marks]			
[Turn over]			



0 7 . 2					
State whether the forward reaction in this equilibrium results in an increase, decrease or no change in the amount, in moles, of gas.					
Explain your answer. [3 marks]					
Tick (✓) ONE box.					
increase					
decrease					
no change					
Explanation					



07.3
Explain why using a catalyst has no effect on the percentage yield. [1 mark]



Hydrogen and nitrogen react to form ammonia.

$$3H_2(g) + N_2(g) \rightleftharpoons 2NH_3(g)$$

At 745 K, the equilibrium constant, $K_c = 0.118 \text{ mol}^{-2} \text{ dm}^6$

At 745 K, 0.150 dm³ of an equilibrium mixture contains 0.0285 mol of hydrogen and 0.0870 mol of nitrogen.

Calculate the amount, in moles, of ammonia present in this equilibrium mixture. [5 marks]

Amount of ammonia ____ mol



			1
0	7		5
_	•	-	•

Calculate the value, at 745 K, for the equilibrium constant $K_{\rm C}$ for this dissociation of ammonia to give hydrogen and nitrogen.

State the units.

$$2 \text{ NH}_3(g) \rightleftharpoons 3 \text{ H}_2(g) + \text{ N}_2(g)$$
 [2 marks]

Units ____

[Turn over]

13



0 8

A student does two test-tube reactions on four colourless solutions (A, B, C and D).

TABLE 4 shows the student's observations.

TABLE 4

SOLUTION	TEST 1	TEST 2
	Add Na ₂ CO ₃ (s)	Add acidified AgNO ₃ (aq)
Α	Effervescence	No visible change
В	Effervescence	White precipitate
С	No visible change	No visible change
D	No visible change	Very pale yellow precipitate



08.1
Identify the gas formed in TEST 1.
Describe a further test to confirm the identity of this gas. [2 marks]
Identity of gas
Test
[Turn over]



BLANK PAGE



0 8 . 2
Explain how the observations from TEST 1 and TEST 2, on page 34, can be used to show that solution B contains hydrochloric acid. [2 marks]



REPEAT OF TABLE 4

SOLUTION	TEST 1	TEST 2
	Add Na ₂ CO ₃ (s)	Add acidified AgNO ₃ (aq)
Α	Effervescence	No visible change
В	Effervescence	White precipitate
С	No visible change	No visible change
D	No visible change	Very pale yellow precipitate

0 0 1.1 3

Describe a series of tests that the student	can use to
show that solution C contains ammonium	sulfate.
[4 marks]	





0 8 . 4
The student does an additional experiment to show that solution D contains a mixture of halide ions. One of the halide ions is chloride.
Method:

Step 1	Add an excess of AgNO ₃ (aq) to 10.0 cm ³ of solution D.
Step 2	Filter, wash, dry and weigh the precipitate.
Step 3	Add an excess of dilute ammonia to the dry precipitate.
Step 4	Filter, wash, dry and weigh the solid that

Explain how the masses recorded during this

remains.

experiment can be used to show that solution D contains a mixture of halide ions. [2 marks]



[Turn over]	10



SECTION B

Answer ALL questions in this section.

Only ONE answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS










If you want to change your answer you must cross out your original answer as shown.



If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.



You may do your working in the blank space around each question but this will not be marked. Do NOT use additional sheets for this working.



Which atom has two more protons and two more neutrons than $^{52}_{24}\text{Cr}$? [1 mark]



A 54/26Cr



B 56/26Cr



C 54₂₆Fe



D 56 26Fe



1 0	
An ato levels.	m has all its electrons in their lowest energy
Which [1 mar	atom contains only two unpaired electrons? k]
0	A Helium
0	B Beryllium
0	C Oxygen
0	D Iron



1	1

The first six ionisation energies, in kJ mol⁻¹, of an element are:

1090, 2350, 4610, 6220, 37 800, 47 000

What is the element? [1 mark]

O A	Boron
-----	-------

O B Carbor





In which pair is the first ionisation energy of atom Y greater than that of atom X? [1 mark]

		Electron configuration of atom X	Electron configuration of atom Y
0	A	1s ² 2s ²	1s ² 2s ² 2p ¹
0	В	1s ² 2s ² 2p ³	1s ² 2s ² 2p ⁴
0	С	1s ² 2s ² 2p ⁵	1s ² 2s ² 2p ⁶
0	D	1s ² 2s ² 2p ⁶	1s ² 2s ² 2p ⁶ 3s ¹




1 3	
	statement about isotopes of an element is NOT t? [1 mark]
0	A They have the same chemical properties.
0	B They have the same number of electrons in ions of the same charge.
0	C They have the same number of neutrons.
0	D They have the same number of protons.



5.0 g of an oxide contains 4.0 g of molybdenum.

What is the empirical formula of this oxide? [1 mark]











1	_
1	J

The equation for a reaction is

$$AsH_3 + H^+ \longrightarrow AsH_4^+$$

What type of interaction forms in this reaction? [1 mark]

- 0
- A Co-ordinate bond
- 0
- B Dipole-dipole force
- 0
- C Hydrogen bond
- 0
- **D** Ionic bond



1 6		
		a correct trend down Group 7 from fluorine to 1 mark]
0	A	The boiling point of the element decreases.
0	В	The oxidising ability of the element decreases.
0	С	The electronegativity of the atom increases.
0	D	The first ionisation energy of the atom increases.



1	7

Which of these ions has the largest ionic radius? [1 mark]











18		
Which	statement	is correct? [1 mark]
0		le ions reduce concentrated sulfuric form sulfur dioxide.
0		le ions reduce concentrated sulfuric form sulfur.
0	C Bromic ions.	le ions reduce iodine to form iodide
0	D lodide ions.	ions reduce chlorine to form chloride



In which of these substances is oxygen in the highest oxidation state? [1 mark]



A OF₂



 BH_2O



 CO_2



 $D H_2O_2$



Which block in the Periodic Table contains the element samarium (Sm)? [1 mark]







O D s block




2	1

Which species is NOT a possible product of the reactions between chlorine and water? [1 mark]











2 2			
Which statement is correct? [1 mark]			
0	A	Magnesium reacts with steam to give magnesium oxide as one of the products.	
0	В	Magnesium acts as an oxidising agent in the extraction of titanium.	
0	С	Magnesium has a lower melting point than sodium.	
0	D	Magnesium hydroxide is very soluble in water.	



2 3			
Which [1 mar		NOT responsible for conducting electricit	:y?
0	A	The sodium ions in molten sodium chlor	'ide
0	В	The electrons between layers of carbon atoms in graphite	
0	С	The bonding electrons in a metal	
0	D	The lone pair electrons in liquid water molecules	
END C)F	QUESTIONS	15



Additional page, if required. Write the question numbers in the left-hand margin.				



Additional page, if required. Write the question numbers in the left-hand margin		



BLANK PAGE

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
Section B		
TOTAL		

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2022 AQA and its licensors. All rights reserved.

IB/M/NC/Jun22/7404/1/E1



