

A



**Surname** \_\_\_\_\_

**Other Names** \_\_\_\_\_

**Centre Number** \_\_\_\_\_

**Candidate Number** \_\_\_\_\_

**Candidate Signature** \_\_\_\_\_

**I declare this is my own work.**

**A-level**

**CHEMISTRY**

**Paper 2 Organic and Physical Chemistry**

**7405/2**

**Time allowed: 2 hours**

**At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.**

**[Turn over]**



J U N 2 2 7 4 0 5 2 0 1

**For this paper you must have:**

- **the Periodic Table/Data Booklet, provided as an insert (enclosed)**
- **a ruler with millimetre measurements**
- **a scientific calculator, which you are expected to use where appropriate.**

## **INSTRUCTIONS**

- **Use black ink or black ball-point pen.**
- **Answer ALL questions.**
- **You must answer the questions in the spaces provided. Do NOT write on blank pages.**
- **If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).**
- **All working must be shown.**



- **Do all rough work in this book. Cross through any work you do not want to be marked.**

## **INFORMATION**

- **The marks for questions are shown in brackets.**
- **The maximum mark for this paper is 105.**


**DO NOT TURN OVER UNTIL TOLD TO DO SO**



**Answer ALL questions in the spaces provided.**

**0 1**

**An acidified solution of butanone reacts with iodine as shown.**



**0 1 . 1**

**On the opposite page, draw the displayed formula for  $\text{CH}_3\text{CH}_2\text{COCH}_2\text{I}$**

**Give the name of  $\text{CH}_3\text{CH}_2\text{COCH}_2\text{I}$**

**[2 marks]**




**Displayed formula**

**Name** \_\_\_\_\_

**[Turn over]**





0 1 . 2

The rate equation for the reaction is

$$\text{rate} = k[\text{CH}_3\text{CH}_2\text{COCH}_3][\text{H}^+]$$

TABLE 1 shows the initial concentrations used in an experiment.

6

TABLE 1

	$\text{CH}_3\text{CH}_2\text{COCH}_3$	$\text{I}_2$	$\text{H}^+$
Initial concentration / $\text{mol dm}^{-3}$	4.35	0.00500	0.825

The initial rate of reaction in this experiment is  
 $1.45 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$




**Calculate the value of the rate constant,  $k$ , for the reaction and give its units. [3 marks]**

**$k$**  \_\_\_\_\_

**Units** \_\_\_\_\_

**[Turn over]**




0 1 . 3

**Calculate the initial rate of reaction when all of the initial concentrations are halved. [1 mark]**

**Initial rate of reaction \_\_\_\_\_ mol dm<sup>-3</sup> s<sup>-1</sup>**





0 9

**BLANK PAGE**

**[Turn over]**

0	1	.	4
---	---	---	---

**An experiment was done to measure the time,  $t$ , taken for a solution of iodine to react completely when added to an excess of an acidified solution of butanone.**

**Suggest an observation used to judge when all the iodine had reacted. [1 mark]**

---

---

---




**The experiment was repeated at different temperatures.**

**FIGURE 1, on page 12, shows how  $\frac{1}{t}$  varied with temperature for these experiments.**

**[Turn over]**



FIGURE 1



0	1	.	5
---	---	---	---

**Describe and explain the shape of the graph in FIGURE 1. [3 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



0	1	.	6
---	---	---	---

**Deduce the time taken for the reaction at 35 °C [1 mark]**

**Time** \_\_\_\_\_ **s**



**BLANK PAGE**

**[Turn over]**






01.7

For a different reaction, TABLE 2 shows the value of the rate constant at different temperatures.

TABLE 2

EXPERIMENT	TEMPERATURE / K	RATE CONSTANT / s <sup>-1</sup>
1	$T_1 = 303$	$k_1 = 1.55 \times 10^{-5}$
2	$T_2 = 333$	$k_2 = 1.70 \times 10^{-4}$



  
This equation can be used to calculate the activation energy,  $E_a$

$$\ln \left( \frac{k_1}{k_2} \right) = \frac{E_a}{R} \left( \frac{1}{T_2} - \frac{1}{T_1} \right)$$

Calculate the value, in  $\text{kJ mol}^{-1}$ , of the activation energy,  $E_a$

The gas constant,  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$  [5 marks]

17

[Turn over]



$E_a$  \_\_\_\_\_  $\text{kJ mol}^{-1}$

18

0 1 . 8

**Name and outline the mechanism for the reaction of butanone with KCN followed by dilute acid. [5 marks]**

**Name of mechanism** \_\_\_\_\_



1 9

# Outline of mechanism


19

[Turn over]

21
----

0	2
---	---

**Tetrafluoroethene is made from chlorodifluoromethane in this reversible reaction.**



**A 2.00 mol sample of  $\text{CHClF}_2$  is placed in a container of volume  $23.2 \text{ dm}^3$  and heated.**

**When equilibrium is reached, the mixture contains  $0.270 \text{ mol}$  of  $\text{CHClF}_2$**



**02.1**

**Calculate the amount, in moles, of  $C_2F_4$  and of HCl in the equilibrium mixture.  
[2 marks]**

**Amount of  $C_2F_4$  \_\_\_\_\_ mol**

**Amount of HCl \_\_\_\_\_ mol**

**[Turn over]**



**02.2**

**Give an expression for  $K_c$  for this equilibrium. [1 mark]**

**$K_c$**



0	2	.	3
---	---	---	---

Calculate a value for  $K_c$

Give its units. [3 marks]

$K_c$  \_\_\_\_\_ Units \_\_\_\_\_

[Turn over]



0	2	.	4
---	---	---	---

**State and explain the effect of using a higher temperature on the equilibrium yield of tetrafluoroethene. [3 marks]**

**Effect on yield** \_\_\_\_\_

**Explanation** \_\_\_\_\_

---

---

---

---

---

---

---

---

---

---





**BLANK PAGE**

**[Turn over]**



**0 2 . 5**

**Chemists provided evidence that was used to support a ban on the use of chlorodifluoromethane as a refrigerant.**

**Many refrigerators now use pentane as a refrigerant.**

**State the environmental problem that chlorodifluoromethane can cause.**

**Give ONE reason why pentane does not cause this problem. [2 marks]**

**Environmental problem** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**Reason why pentane does not cause this problem**

\_\_\_\_\_

\_\_\_\_\_



---

---

---

---

---

**[Turn over]**

<b>11</b>



0	3
---	---

**This question is about 2-methylbut-1-ene.**

0	3	.	1
---	---	---	---

**Name the mechanism for the reaction of 2-methylbut-1-ene with concentrated sulfuric acid.**

**Outline the mechanism for this reaction to form the major product. [5 marks]**

**Name of mechanism** \_\_\_\_\_

---

**Outline of mechanism to form major product**



**[Turn over]**



03.2

**Draw the structure of the minor product formed in the reaction in Question 03.1**

**Explain why this is the minor product.  
[3 marks]**

**Structure of minor product**

**Explanation** \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_



0	3	.	3
---	---	---	---

**Draw the skeletal formula of a functional group isomer of 2-methylbut-1-ene.  
[1 mark]**

**[Turn over]**



0	3	.	4
---	---	---	---

**2-methylbut-1-ene can form a polymer.**

**State the type of polymerisation.**

**Draw the repeating unit for the polymer formed. [2 marks]**

**Type of polymerisation** \_\_\_\_\_

---

**Repeating unit**





**BLANK PAGE**

**[Turn over]**



0	4
---	---

**Proteins are polymers made from amino acids.**

**Part of the structure of a protein is shown.**

**–Cys–Ser–Asp–Phe–**

**Each amino acid in the protein is shown using the first three letters of its name.**

0	4	.	1
---	---	---	---

**Identify the type of protein structure shown. [1 mark]**

**Tick (✓) ONE box.**

**Primary**

**Secondary**

**Tertiary**



**04.2**

**Draw a structure for the –Cys–Ser–  
section of the protein.**

**Use the Data Booklet to help you answer  
this question. [2 marks]**

**04.3**


**Name the other substance formed when  
two amino acids react together to form  
part of a protein chain. [1 mark]**

---

**[Turn over]**



The general structure of an amino acid is shown.



R represents a group that varies between different amino acids.

R groups can interact and contribute to protein structure.

04.4

Explain why the strength of the interaction between two cysteine R groups differs from the strength of the interaction between a serine R group and an aspartic acid R group.

Use the Data Booklet to help you answer this question. [4 marks]





0	4	.	5
---	---	---	---

**Deduce the type of interaction that occurs between a lysine R group and an aspartic acid R group. [1 mark]**

---

9



**BLANK PAGE**

**[Turn over]**



0	5
---	---

**This question is about the preparation of hexan-2-ol.**

**Hexan-2-ol does not mix with water and has a boiling point of 140 °C**

**Hexan-2-ol can be prepared from hex-1-ene using this method.**

- A Measure out 11.0 cm<sup>3</sup> of hex-1-ene into a boiling tube in an ice bath.**
- B Carefully add 5 cm<sup>3</sup> of concentrated phosphoric acid to the hex-1-ene.**
- C After 5 minutes add 10 cm<sup>3</sup> of distilled water to the mixture and transfer the boiling tube contents to a separating funnel.**
- D Shake the mixture and allow it to settle.**
- E Discard the lower (aqueous) layer.**





- F** Add a fresh 10 cm<sup>3</sup> sample of distilled water and repeat steps D and E.
- G** Transfer the remaining liquid to a beaker.
- H** Add 2 g of anhydrous magnesium sulfate and allow to stand for 5 minutes.
- I** Filter the mixture under reduced pressure.
- J** Distil the filtrate and collect the distillate that boils in the range 130–160 °C

**[Turn over]**



**BLANK PAGE**



**05.1**

**It is important to wear eye protection and a lab coat when completing this experiment.**

**Suggest, with a reason, ONE other appropriate safety precaution for this experiment. [2 marks]**

**Precaution** \_\_\_\_\_

\_\_\_\_\_

**Reason** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**[Turn over]**



**05.2**

**Give a reason for adding the distilled water in steps C and F. [1 mark]**

---

---

---

**05.3**

**Give a reason for adding anhydrous magnesium sulfate in step H. [1 mark]**

---


---

---



**0 5 . 4**

**Complete and label the diagram of the apparatus used to filter the mixture under reduced pressure in step 1. [2 marks]**



**[Turn over]**



**0 5 . 5**

**Identify the most likely organic impurity, other than hex-1-ene, in the distillate collected in step J.**

**Suggest ONE reason why it could be difficult to remove this impurity. [2 marks]**

**Impurity** \_\_\_\_\_

**Reason** \_\_\_\_\_

**0 5 . 6**

**On the opposite page, calculate the mass, in g, of hexan-2-ol formed from 11.0 cm<sup>3</sup> of hex-1-ene if the yield is 31.0%**

**Give your answer to 1 decimal place.**

**Density of hex-1-ene = 0.678 g cm<sup>-3</sup>  
[4 marks]**



Mass \_\_\_\_\_ g

[Turn over]



4 7

12

0	6
---	---

**This question is about compound X with the empirical formula  $C_2H_4O$**

**FIGURE 2, on page 50, shows the infrared spectrum of X.**

**FIGURE 3, on page 51, shows the  $^{13}C$  NMR spectrum of X.**

**The  $^1H$  NMR spectrum of X shows four peaks with different chemical shift values.**

**TABLE 3, on page 51, gives data for these peaks.**





**BLANK PAGE**

**[Turn over]**



## FIGURE 2

Transmittance / %




FIGURE 3




TABLE 3

<b>Chemical shift <math>\delta</math> / ppm</b>	<b>3.9</b>	<b>3.7</b>	<b>2.1</b>	<b>1.2</b>
<b>Splitting pattern</b>	<b>quartet</b>	<b>singlet</b>	<b>singlet</b>	<b>doublet</b>
<b>Integration value</b>	<b>1</b>	<b>1</b>	<b>3</b>	<b>3</b>



**BLANK PAGE**





Lined writing area with 18 horizontal lines.



---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**[Turn over]**



5 7

**6**



07


**This question is about esters.**

**FIGURE 4, on the opposite page, shows an incomplete mechanism for the reaction of an ester with aqueous sodium hydroxide.**

07.1

**Add THREE curly arrows to complete the mechanism in FIGURE 4. [3 marks]**

**FIGURE 4**



[Turn over]



07.2

**Name the type of reaction shown in FIGURE 4, on page 59.**  
**[1 mark]**

---

---

**60**

07.3

**Deduce the role of the  $\text{CH}_3\text{O}^-$  ion in step 3 shown in FIGURE 4, on page 59. [1 mark]**

---

---



07.4

**A triester in vegetable oil reacts with sodium hydroxide in a similar way.**

**Give a use for a product of this reaction. [1 mark]**

---

**61**

---

**[Turn over]**

**6**

0	8
---	---

**Benzene reacts with methanoyl chloride (HCOCl) in the presence of a catalyst.**

0	8	.	1
---	---	---	---

**Give an equation for the overall reaction when benzene reacts with methanoyl chloride.**

**Name the organic product. [2 marks]**

**Equation** \_\_\_\_\_

**Name** \_\_\_\_\_



0	8	.	2
---	---	---	---

**Identify the catalyst needed in this reaction.**

**Give an equation to show how the catalyst is used to form the electrophile,  $[\text{HCO}]^+$**

**[2 marks]**

**Catalyst** \_\_\_\_\_

\_\_\_\_\_

**Equation** \_\_\_\_\_

**[Turn over]**



0	8	.	3
---	---	---	---

**Outline the mechanism for the reaction of benzene with the electrophile,  $[\text{HCO}]^+$   
[3 marks]**

7





**BLANK PAGE**

**[Turn over]**





0	9
---	---


**This question is about olive oil.**

**A sample of olive oil is mainly the unsaturated fat Y mixed with a small amount of inert impurity.**

**The structure of Y in the olive oil is shown.**

**Y has the molecular formula  $C_{57}H_{100}O_6$  ( $M_r = 880$ ).**

**66**





**The amount of Y is found by measuring how much bromine water is decolourised by a sample of oil, using this method.**

- Transfer a weighed sample of oil to a 250 cm<sup>3</sup> volumetric flask and make up to the mark with an inert organic solvent.**
- Titrate 25.0 cm<sup>3</sup> samples of the olive oil solution with 0.025 mol dm<sup>-3</sup> Br<sub>2</sub>(aq).**

**67**

**[Turn over]**



09.1

A suitable target titre for the titration is  $30.0 \text{ cm}^3$  of  $0.025 \text{ mol dm}^{-3} \text{ Br}_2(\text{aq})$ .

Justify why a much smaller target titre would NOT be appropriate.

Calculate the amount, in moles, of bromine in the target titre. [2 marks]

68

Justification

---

---

---

---

---

---

---



Amount of bromine \_\_\_\_\_ mol

**[Turn over]**

**09.2**

**Calculate a suitable mass of olive oil to transfer to the volumetric flask using your answer to Question 09.1 and the structure of Y.**

**Assume that the olive oil contains 85% of Y by mass.**

**(If you were unable to calculate the amount of bromine in the target titre, you should assume it is  $6.25 \times 10^{-4}$  mol. This is NOT the correct amount.) [5 marks]**



Mass of olive oil \_\_\_\_\_ g

[Turn over]



**The olive oil solution can be prepared using this method.**

- **Place a weighing bottle on a balance and record the mass, in g, to 2 decimal places.**
- **Add olive oil to the weighing bottle until a suitable mass has been added.**
- **Record the mass of the weighing bottle and olive oil.**
- **Pour the olive oil into a 250 cm<sup>3</sup> volumetric flask.**
- **Add organic solvent to the volumetric flask until it is made up to the mark.**
- **Place a stopper in the flask and invert the flask several times.**





09.3

**Suggest an extra step to ensure that the mass of olive oil in the solution is recorded accurately.**

**Justify your suggestion. [2 marks]**

**Extra step** \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**Justification** \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**[Turn over]**



0	9	.	4
---	---	---	---

**State the reason for inverting the flask several times. [1 mark]**

---

---

---

---

---

---



**BLANK PAGE**

**[Turn over]**



**09.5**

**A sample of the olive oil was dissolved in methanol and placed in a mass spectrometer. The sample was ionised using electrospray ionisation. Each molecule gained a hydrogen ion (H<sup>+</sup>) during ionisation.**

**The spectrum showed a peak for an ion with  $\frac{m}{z} = 345$  formed from an impurity in the olive oil.**

**The ion with  $\frac{m}{z} = 345$  was formed from a compound with the empirical formula  $C_5H_{10}O$**

**Deduce the molecular formula of this compound. [2 marks]**

**Show your working.**



**Molecular formula** \_\_\_\_\_

**[Turn over]**



<b>12</b>



1 0

**This question is about the reaction scheme shown on the opposite page.**

1 0 . 1

**State the reagents needed for step 1 and the reagents needed for step 2. [3 marks]**

**78**

**step 1**

---

---

---

**step 2**





---

---

---


---

---



[Turn over]

**BLANK PAGE**







8 1

1 0 . 2

**Give the name of the mechanism for the reaction in step 3.**  
**[1 mark]**

---

1 0 . 3

**81**

**Name the reagent for step 4.**

**State a necessary condition for step 4. [2 marks]**

**Reagent** \_\_\_\_\_

**Condition** \_\_\_\_\_

**[Turn over]**



1 0 . 4

**Amine A is formed in step 2 and amine B is formed in step 5.**

**Explain why the yield of B in step 5 is less than the yield of A in step 2. [2 marks]**

---

---

---

---

---

---

---



1 0 . 5

**Explain why amine B is a stronger base than amine A.**

**[2 marks]**

---

---

---

---

---

---

---

**83**

**10**

**END OF QUESTIONS**



**Additional page, if required.**  
**Write the question numbers in the left-hand margin.**


**BLANK PAGE**

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
<b>TOTAL</b>	

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk).

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2022 AQA and its licensors. All rights reserved.

**IB/M/SB/Jun22/7405/2/E2**

8 6



2 2 6 A 7 4 0 5 / 2