

A



# **Physics Equations Sheet**

## **GCSE Physics (8463)**

### **FOR USE IN JUNE 2022 ONLY**

**[Turn over]**

## HT = Higher Tier only equations

kinetic energy = $0.5 \times \text{mass} \times (\text{speed})^2$	$E_k = \frac{1}{2} m v^2$
elastic potential energy = $0.5 \times \text{spring constant} \times (\text{extension})^2$	$E_e = \frac{1}{2} k e^2$
gravitational potential energy = mass $\times$ gravitational field strength $\times$ height	$E_p = m g h$
Change in thermal energy = mass $\times$ specific heat capacity $\times$ temperature change	$\Delta E = m c \Delta \theta$
power = $\frac{\text{energy transferred}}{\text{time}}$	$P = \frac{E}{t}$

<b>power = <math>\frac{\text{work done}}{\text{time}}</math></b>	<b><math>P = \frac{W}{t}</math></b>
<b>efficiency = <math>\frac{\text{useful output energy transfer}}{\text{total input energy transfer}}</math></b>	
<b>efficiency = <math>\frac{\text{useful power output}}{\text{total power input}}</math></b>	
<b>charge flow = current × time</b>	<b><math>Q = I t</math></b>
<b>potential difference = current × resistance</b>	<b><math>V = I R</math></b>
<b>power = potential difference × current</b>	<b><math>P = V I</math></b>

<b>power = (current)<sup>2</sup> × resistance</b>	$P = I^2 R$
<b>energy transferred = power × time</b>	$E = P t$
<b>energy transferred = charge flow × potential difference</b>	$E = Q V$
<b>density = <math>\frac{\text{mass}}{\text{volume}}</math></b>	$\rho = \frac{m}{V}$
<b>Thermal energy for a change of state = mass × specific latent heat</b>	$E = m L$
<b>For gases: pressure × volume = constant</b>	$p V = \text{constant}$
<b>weight = mass × gravitational field strength</b>	$W = m g$
<b>work done = force × distance (along the line of action of the force)</b>	$W = F s$

<b>force = spring constant × extension</b>	$F = k e$
<b>moment of a force = force × distance (normal to direction of force)</b>	$M = F d$
<b>pressure = <math>\frac{\text{force normal to a surface}}{\text{area of that surface}}</math></b>	$p = \frac{F}{A}$
<b>Pressure due to column of liquid = height of column × density of liquid × gravitational field strength</b>	$p = h \rho g$
<b>distance travelled = speed × time</b>	$s = v t$
<b>acceleration = <math>\frac{\text{change in velocity}}{\text{time taken}}</math></b>	$a = \frac{\Delta v}{t}$

HT

[Turn over]



<p><b>HT</b></p> <p><b>Force on a conductor (at right angles to a magnetic field) carrying a current = magnetic flux density × current × length</b></p>	$F = B I l$
<p><b>HT</b></p> <p><b><u>potential difference across primary coil</u></b>  <b><u>potential difference across secondary coil</u></b>  <b>= <u>number of turns in primary coil</u></b>  <b><u>number of turns in secondary coil</u></b></p>	$\frac{V_p}{V_s} = \frac{n_p}{n_s}$
<p><b>HT</b></p> <p><b>potential difference across primary coil × current in primary coil = potential difference across secondary coil × current in secondary coil</b></p>	$V_p I_p = V_s I_s$

**BLANK PAGE**

**Physics Equations Sheet - GCSE  
Physics (8463)**

**FOR USE IN JUNE 2022 ONLY**

**IB/M/CD/Jun22/8463/INS/E1**