

Surname	
Forename(s)	
Centre Number	
Candidate Number	
Candidate Signature	
I declare this is my own work.	

AS BIOLOGY

Paper 1

7401/1

Monday 15 May 2023 Morning

Time allowed: 1 hour 30 minutes



At the front of this book, write your surname and forename(s), your centre number, your candidate number and add your signature.

MATERIALS

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Answer ALL questions.
- You must answer the questions in the spaces provided. Do not write on blank pages.



- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

INFORMATION

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 75.

DO NOT TURN OVER UNTIL TOLD TO DO SO



Answer ALL questions in the spaces provided.

0 1.1 Describe the primary structure of all					
	[2 marks]				



0 1 . 2	0	1		2
---------------	---	---	--	---

This question is about the genetic code.

Define UNIVERSAL, NON-OVERLAPPING and DEGENERATE. [3 marks]


Universal		
Non-overlapping _		
Degenerate		



0 2

FIGURE 1 shows a transmission electron microscope (TEM) image of a 'Streptococcus' bacterium.

FIGURE 1





0	2	1

Describe how the appearance of the area containing DNA in a TEM image of a eukaryotic cell would differ from that shown in FIGURE 1. [2 marks]



0 2 .

Describe ONE difference between the structure of DNA in a prokaryotic cell and in a eukaryotic cell. [1 mark]

- - -

'Streptococcus' bacteria can infect the lungs when air is breathed in and cause lung disease.

Describe the mechanism of breathing that causes air to enter the lungs.
[3 marks]





Some strains of 'Streptococcus' bacteria are more likely to cause lung disease than other strains.

Strains that do not cause lung disease are quickly destroyed by phagocytes. Phagocytes are stimulated when they bind to murein on 'Streptococcus' bacteria.

Each strain of 'Streptococcus' bacteria has a capsule of different thickness from the others.

Suggest how 'Streptococcus' bacteria with a thicker capsule are more likely to survive AND so cause lung disease.
[2 marks]



[Turn over]	8







FIGURE 2 shows a fatty acid that contains 20 carbon atoms and four double bonds.

FIGURE 2



On FIGURE 2, draw a box around the R group of the fatty acid. [1 mark]




BLANK PAGE



03.2

FIGURE 3 shows a triglyceride.

FIGURE 3



KEY

Glycerol

/// Ester bond

Fatty acid



Describe TWO differences betw	veen the
structure of the triglyceride she	own in
FIGURE 3 and a phospholipid.	[2 marks]

1			
2			



0	3		3
		_	


Explain why phospholipids can form a bilayer but triglycerides cannot. [3 marks]




BLANK PAGE



hows two fatty acids, A and B. 0 3 . 4 FIGURE 4 sh FIGURE 4





ed rats a diet with added fish oil for 4 months. Scientists for

before starting this diet (0 months) and after 4 months on They obtained samples of red blood cells from the rats this diet.

d blood cell sample, they separated the membranes and measured: cell-surface For each re

ntage of phospholipids containing each of the fatty acids A and B the perce


• the fluidity of the membrane.

page 20, shows the scientists' results. TABLE 1, on

Time	Mean	Mean	Mean
sample of	percentage of	percentage of	fluidity of
red blood	phospholipids	phospholipids	the
cells	containing	containing fatty	membrane /
obtained /	fatty acid A	acid B	arbitrary
months			units
0	19.8	1.7	31
4	11.7	0'6	26



Suggest why the fluidity of the membrane was higher after 4 months.
Use all the information provided in the question. [3 marks]





					9



BLANK PAGE



0 4

Scientists investigated the activation of T lymphocytes.

The scientists studied two types of cell:

- naïve T cells, which are T cells that have not yet been in contact with a foreign antigen
- activated helper T cells, which are T cells that have been activated by a foreign antigen.


FIGURE 5, on the opposite page, shows electron microscope images of the two types of cell.




FIGURE 5

NAÏVE T CELL

ACTIVATED HELPER T CELL





The activated helper T cell has a volume of 463 μm^3



04.1

Calculate the volume of the naïve T cell shown in FIGURE 5.

FIGURE 5 is provided on page 25.

Then calculate how many times larger the activated helper T cell volume is compared with the naïve T cell volume.

Assume the cell is spherical.

Use line PQ to measure the diameter of the naïve T cell.

Volume of a sphere $=\frac{4}{3}\pi r^3$ where π is 3.14

Show your working, on the opposite page. [3 marks]



Volume of naïve	T cell	μm ³

Number of times larger the activated helper T cell volume is compared with the naïve T cell volume



0	4	•	2

State ONE feature that shows the images in FIGURE 5, on page 25, were taken with an electron microscope and NOT an optical microscope.

Explain	ı your	ansv	ver. [2 mar	'KS]	



|--|

State ONE role of a helper T cell. [1 mark]



The rate of translation is increased in T cells activated by antigens.

Describe the role of tRNA in translation. [2 marks]





BLANK PAGE



0 5


A student investigated the effect of changing surface area on the rate of diffusion of a solution into the centre of agar blocks.


She used agar coloured by an indicator. The indicator is pink at pH > 8 and colourless at pH 8 and pH < 8

She cut blocks in different shapes as shown in FIGURE 6, on the opposite page.



FIGURE 6







BLANK PAGE




0 5 . 1

Complete TABLE 2 to show the surface area and the surface area to volume ratio for the two shapes, on page 33.
[2 marks]

TABLE 2

Shape	Volume / cm ³	Surface area to volume ratio
С	64	:1
D	64	:1





put the blocks into an acidic solution. The studen The acidic solution caused the blocks to gradually turn from pink to colourless.

colourless. She repeated this three times. She recorded the time taken for the blocks to turn completely

n the opposite page, shows the student's TABLE 3, o results


TABLE 3

Time for / s	ime for block to turn colous	urn colot	ırless	Mean time for	
Block 1	Block 2	Block 3	Block 4	colourless / s	
3490	1200	3540	3530		
 1680	1500	1590	1610	1595	3 <i>1</i>

After collecting the data, the student noticed that shape C, damaged. block 2 was

le mean for shape C. [1 mark] Calculate th







3 9

State THREE variables the student controlled in order to obtain valid results. [2 marks] results. [2 marks]

0	5		5
		_	

bescribe now gas exchange occurs in single-celled organisms AND explain why this method cannot be used by large, multicellular organisms. [3 marks	. 1
iargo, marcioonalar organionior jo marko	



Γ	Λ	6
	U	O

The human papilloma virus contains a double-stranded DNA genome.

0	6		1
---	---	--	---

Which components are found in a human papilloma virus? [1 mark]

Tick (✓) ONE box.

Capsid, attachment protein and
reverse transcriptase

	Capsule and attachment	protein
--	------------------------	---------

Cell-surface membrane and
attachment protein



0	6		2
		-	

The DNA-replication enzymes of a human cell make copies of the human papilloma virus genome.

Name TWO enzymes that are involved in replicating the DNA of the human papilloma virus and describe their roles in the replication process. [3 marks]

Name of enzyme 1	
Role of enzyme 1	
Name of enzyme 2	



Role of enzyme 2		_



0	6	3
_	_	 _

Human papilloma virus infects cells that are no longer dividing. The human papilloma virus genome contains genes that code for proteins that cause human cells to restart their cell cycles.

Human papilloma virus infection can cause cancer.

Explain wny. [1 mark]			





06.4

Human papilloma virus (HPV) is transmitted through sexual contact.

More than 95% of cervical cancers (which only affect females) are due to HPV infection. HPV infection of other tissues increases the risk of cancer but this is rare compared with cervical infection.

A vaccine is available that is over 80% effective at preventing HPV infection, if given before the person has been exposed to HPV. There is evidence of herd immunity when more than 50% of the population have been vaccinated.

Evaluate whether 10- to 12-year-old boys should be given the HPV vaccine.
[4 marks]



[Turn over]	9

4 7

0 7

Scientists investigated biodiversity in prokaryote communities found in soil.

The scientists:

- took soil samples from fields that had been managed for 20 years with two different farming methods
- sequenced all the DNA that coded for prokaryotic ribosomal RNA in the soil samples
- compared these base sequences to give a measure of species richness and an index of diversity for the prokaryote community
- recorded the total prokaryotic biomass and the mass of stored carbon for each soil sample
- obtained the mean wheat yield from the fields.





TABLE 4, below and on the opposite page, shows the scientists' results.

TABLE 4

DATA COLLECTED	FARMING METHOD 1	FARMING METHOD 2
Mean species richness (± 2 × standard deviation)	517 (± 17)	560 (± 24)
Mean index of diversity (± 2 × standard deviation)	0.251 (± 0.011)	0.230 (± 0.014)
Mean total prokaryotic biomass / kg m ⁻³	0.24	0.40



DATA COLLECTED	FARMING METHOD 1	FARMING METHOD 2
Mean carbon stored in soil organisms / µg g ⁻¹	203	342
Mean wheat yield / g m ⁻²	451	377

The mean ± 2 × standard deviation includes 95% of the data.





0	7	1
		•

Using the standard deviation data from TABLE 4, on pages 50 and 51, describe the differences in prokaryotic biodiversity found in the soil with these two farming methods.

In your answer, give the definitions of

SPECIES RICHNESS and INDEX OF DIVERSITY. [4 marks]			







0	7	2
		_

Genetic diversity in soil species was traditionally inferred by making observations after growing prokaryotes on agar plates.

However, it is estimated that less than 10% of prokaryotes found in soil will grow if spread on an agar plate in a laboratory.

In recent years, our knowledge of prokaryotic biodiversity in the soil has increased.

Suggest why. [2 marks]			



Cuara at white



REPEAT OF TABLE 4

DATA COLLECTED	FARMING METHOD 1	FARMING METHOD 2
Mean species richness (± 2 × standard deviation)	517 (± 17)	560 (± 24)
Mean index of diversity (± 2 × standard deviation)	0.251 (± 0.011)	0.230 (± 0.014)
Mean total prokaryotic biomass / kg m ⁻³	0.24	0.40



DATA COLLECTED	FARMING METHOD 1	FARMING METHOD 2
Mean carbon stored in soil organisms / µg g ⁻¹	203	342
Mean wheat yield / g m ⁻²	451	377

The mean ± 2 × standard deviation includes 95% of the data.





_		
		7
•	7	7.

Evaluate the balance between conservation and farming for these two farming methods.

Use the information provided in TABLE 4, on pages 58 and 59. [2 mark				

[Turn over]



8

08.1
A student prepared a plant root to observe cells undergoing mitosis.
He put the root in a small bottle of hydrochloric acid in a 40 °C water bath.
Why did he put the plant root in acid? [1 mark]



State TWO precautions required when working with hydrochloric acid. [2 marks]

1			
2			



0	8	3

Scientists determined the mean mitotic index at 50 µm intervals away from the root tip in 10 young plant roots.

FIGURE 7, on the opposite page, shows the scientists' results.

State the null hypothesis for this investigation.


Name the statistical test needed to determine whether the difference between the mean mitotic index at 200 µm and at 300 µm is significant. [2 marks]

Nuii nypotnesis _		
Statistical test		



FIGURE 7


Mean mitotic index





REPEAT OF FIGURE 7

Mean mitotic index



08.4

The scientists recorded these measurements from the tissue located at 200 µm from the root tip.



Area of field of view = 0.2 mm^2 Mean area of one cell = $3495 \mu \text{m}^2$

Use this information and FIGURE 7, on the opposite page, to calculate the number of cells undergoing mitosis in this location.

Assume there are no spaces between the cells.

Show your working. [2 marks]

cells



	0		
U	0	-	J

Another student compares the mitotic index in the roots of two different species.

Give TWO considerations within her method to ensure this comparison is valid. [2 marks]

1			
2			

9





0 9

Read the following passage.

Carbohydrates are moved by mass transport in plants and in mammals. This movement is caused by a pressure gradient inside the transport vessels.

5

Plants transport most of their carbohydrate as sucrose. The phloem has sucrose-transport proteins in the cell-surface membrane. Potato plants with fewer 10 sucrose-transport proteins produce fewer potatoes and accumulate sugars in their leaves.

Mammals do not have sucrose-transport proteins in their 15 cell-surface membranes. They do have many monosaccharide-transport proteins in



cell-surface membranes. These transport proteins allow monosaccharides through by facilitated diffusion.

20

Mammalian cells can change the number of glucose-transport proteins in their cell-surface membranes. The 25 glucose-transport proteins are stored in internal membranes, then moved to the cell-surface membrane when they are needed. During exercise, there is a significant increase in glucose 30 uptake by muscle cells.

Use the information in the passage and your own knowledge to answer the following questions.



Explain how sucrose-transport proteins in leaf cells enable the production of a pressure gradient in the phloem. [3 marks]





0	9	2
		_

Potato plants with fewer sucrose-transport proteins produce fewer potatoes and accumulate sugars in their leaves (lines 10 to 13).

Explain why. [2 marks]					





0	9		3
		_	

During exercise, there is a significant increase in the glucose uptake by muscle cells (lines 29 to 31).

Give two reasons why glucose uptake by muscle cells increases significantly during exercise.


Explain your answers. [4 marks]

1				



2			





Which statement correctly describes the structure of both a sucrose-transport protein and a glucose-transport protein? [1 mark]

Tick (✓) ONE box.

The secondary structure is held by
disulfide bridges. The tertiary
structure allows the protein to be
positioned on the inner surface of
the cell membrane.

The secondary structure is held by disulfide bridges. The tertiary structure allows the protein to be positioned spanning the whole cell membrane.



	The secondary structure is held hydrogen bonds. The tertiary structure allows the protein to be positioned on the outer surface the cell membrane.	е
	The secondary structure is held hydrogen bonds. The tertiary structure allows the protein to be positioned spanning the whole membrane.	е
FND	OF OUESTIONS	10



Additional page, if required. Write the question numbers in the left-hand margin.



Additional page, if required.			
Write the question numbers in the left-hand margin.			



For Examiner's Use			
Question	Mark		
1			
2			
3			
4			
5			
6			
7			
8			
9			
TOTAL			

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 AQA and its licensors. All rights reserved.

WP/M/CD/Jun23/7401/1/E3



