AQA	

Surname

Forename(s)

Centre Number

Candidate Number

Candidate Signature _____

I declare this is my own work.

GCSE CHEMISTRY

Foundation Tier Paper 2 8462/2F

Tuesday 13 June 2023

Morning

Time allowed: 1 hour 45 minutes

At the top of the page, write your surname and forename(s), your centre number, your candidate number and add your signature.



MATERIALS

For this paper you must have:

- a ruler
- a scientific calculator
- the periodic table (enclosed).

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Pencil should only be used for drawing.
- Answer ALL questions in the spaces provided. Do not write on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).



- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

INFORMATION

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

DO NOT TURN OVER UNTIL TOLD TO DO SO



0 1

This question is about oxygen.

Scientists think that there was little or no oxygen in the Earth's early atmosphere.



0	1		1
---	---	--	---

Which planet today has an atmosphere that is similar to the Earth's early atmosphere? [1 mark]


Tick (✓) ONE box.			
	Jupiter		
	Mars		
	Neptune		
	Saturn		



0 1.2

Which is the approximate percentage of oxygen in the Earth's atmosphere today? [1 mark]

Tick (✓) ONE box.



50%

80%

100%



01.3

Which TWO of the following increased the percentage of oxygen in the Earth's atmosphere? [2 marks]

Tick (✓) TWO boxes.

	Active vo	canoes	emitted	gases

Algae and plants evolved

Animals evolved

Carbonate sediments formed in oceans

Photosynthesis took place



01.4

Some scientists think that 1100 million years ago the Earth's atmosphere contained:

- 16% oxygen
- 4% carbon dioxide.

Complete FIGURE 1, on the opposite page.

You should:


- complete the y-axis scale
- plot the percentage of oxygen in the Earth's atmosphere 1100 million years ago.

[2 marks]



FIGURE 1

Percentage (%) of gas in the Earth's atmosphere 1100 million years ago





Oxygen is produced when manganese dioxide is added to hydrogen peroxide solution.

The equation for the reaction is:

hydrogen peroxide --->

water + oxygen

A student investigated the effect of changing the temperature on the decomposition of hydrogen peroxide.


This is the method used.

- 1. Add 5 cm³ of hydrogen peroxide solution to three test tubes labelled A, B and C.
- 2. Place each test tube in a water bath at a different temperature.
- 3. Add 0.2 g of manganese dioxide to each test tube.



FIGURE 2 shows the results.

FIGURE 2





BLANK PAGE



0 1.5

Which test tube contained hydrogen peroxide solution at the highest temperature? [1 mark]

lick	(✓) ONE box.
	Test tube A
	Test tube B
	Test tube C



0	1	•	6
---	---	---	---

The student tested the gas produced.

What is used to prove that the gas is oxygen? [1 mark]

Tick (✓) ONE box.

A glowing splint
Bromine water

I Pailip Illiido papoi		Damp	litmus	paper
------------------------	--	------	--------	-------

01.7

Manganese dioxide does not appear in the chemical equation for this reaction.



Which is a correct statement about manganese dioxide in this reaction? [1 mark]

Tick (√) ONE box.
	Manganese dioxide increases the activation energy in this reaction.
	Manganese dioxide is a catalyst in this reaction.
	Manganese dioxide is used up during this reaction.
	Manganese dioxide reduces the rate of this reaction.
[Turn	over]



0 2

This question is about glass and polymers.

Beakers can be made from borosilicate glass or poly(propene).

TABLE 1, on page 18, shows information about materials used to make beakers.



BLANK PAGE



TABLE 1

	Material used to make beakers		
	borosilicate poly(propene glass		
Temperature at which melting begins in °C	850	160	
Flammability	does not burn	burns	
Resistance to impact	shatters	tough	
Cost of 100 cm ³ beaker in £	1.50	2.00	



		4
0	2	1

Suggest TWO reasons why a Bunsen burner should NOT be used to heat a liquid in a poly(propene) beaker.

Use TABLE 1. [2 marks]				
1				
2				



02.2

Poly(propene) beakers are more expensive than borosilicate glass beakers.

Suggest ONE reason why using poly(propene) beakers instead of borosilicate glass beakers could save money.

Use TABLE 1, on page 18. [1 mark]



Which is a raw material used to make borosilicate glass? [1 mark]

Tick (✓) ONE box.

	Boron	trioxide
--	-------	----------

Clay

Limestone



Poly(propene) is produced from propene.

The displayed structural formula of propene is:

TABLE 2, on the opposite page, shows some information about the elements in one molecule of propene.



TABLE 2

Symbol for element	Name of element	Number of atoms of element in one molecule of propene
C		
Н		


Complete TABLE 2. [2 marks]







Which structure is the repeating unit of poly(propene)? [1 mark]

Tick (✓) ONE box.









BLANK PAGE



02.6

Poly(propene) is produced in three stages:

- STAGE 1: separating large alkane molecules from crude oil
- STAGE 2: producing propene molecules from large alkane molecules
- STAGE 3: joining many propene molecules together.

Name STAGE 1, STAGE 2 and STAGE 3.

Choose answers from the list. [3 marks]

- cracking
- fermentation
- fractional distillation
- polymerisation
- reverse osmosis



STAGE 1 is	-
STAGE 2 is	_
STAGE 3 is	•



0	2		7
		_	_

A molecule of hexene contains a double carbon–carbon bond.

Many hexene molecules join together to form poly(hexene).

Which TWO words describe a hexene molecule in this process? [2 marks]


Tick (✓) TWO boxes.

Alkene
Catalyst
Composite
Element
Monomer



BLANK PAGE






This question is about chromatography.

A student investigated an orange dye using paper aphy. chromatogra

n the opposite page, shows the apparatus at the nvestigation. start of the in FIGURE 3, o

e labels on FIGURE 3. [3 marks] Complete the




[Turn over]



03.2

FIGURE 4 shows the results at the end of the investigation.

FIGURE 4





The student made a mistake in the investigation.

What mistake did the student make to produce the results shown in FIGURE 4? [1 mark]

Tick (✓) ONE box.


Left the investigation for too long
Used a lid on the beaker
Used a solvent which did not dissolve the dye



A different student did the investigation correctly.

FIGURE 5 shows the results.

FIGURE 5



03.3

How do the results in FIGURE 5 show that the orange dye is NOT a pure substance? [1 mark]



BLANK PAGE



0 3 . 4

Determine the R_f value for the red spot.

You should measure:

- the distance moved by the red spot
- the distance moved by the solvent.

Use FIGURE 5, on page 34 and the equation:

R_f = distance moved by red spot distance moved by solvent

[4 marks]

Distance moved by red spot

____cm

Distance moved by solvent

cm



R _f =
03.5
Which spot had the greatest R _f value?
Use FIGURE 5, on page 34. [1 mark]
Tick (✓) ONE box.
Dark yellow spot
Pale yellow spot
Red spot

[Turn over]

0 4

This question is about a reversible reaction.

A student heated calcium hydroxide to produce calcium oxide and water vapour.

This is the method used.

- 1. Add 2.00 g of calcium hydroxide into a test tube.
- 2. Heat the test tube and contents for 1 minute using a Bunsen burner.
- 3. Allow the test tube and contents to cool.
- 4. Weigh the test tube and contents.
- 5. Repeat steps 2 to 4 five more times.



BLANK PAGE



0 4.1

TABLE 3 gives the appearance of the reactant and of the products.

TABLE 3

	COMPOUND	APPEARANCE
REACTANT	calcium hydroxide	white powder
PRODUCTS	calcium oxide	white powder
	water vapour	colourless gas



The student looked at the test tube and contents during heating.

The student could NOT tell that a chemical reaction was taking place by looking at the test tube and contents.

Give TWO reasons why.

Use the information in TABLE 3. [2 marks]		
1		
2		



|--|

Accurate results are NOT produced if solid powders escape from the test tube during heating.

Suggest why sealing the test tube with a stopper is NOT a good way of preventing the solid powders from escaping.
[1 mark]

0	4	•	3

The student wanted to calculate the mass of the contents of the test tube after each minute of heating.



The student weighed the test tube and contents after each minute of heating.

What OTHER measurement is also needed to calculate the mass of the contents of the test tube? [1 mark]

Tick (✓) ONE box.

The change in mass of the

contents of the test tube at the end

The mass of the contents of the test tube at the start

The mass of the empty test tube



The student heated 2.00 g of calcium hydroxide to produce calcium oxide and water vapour.

TABLE 4 shows the results.

TABLE 4

Total heating time in minutes	Mass of contents of test tube in grams
0	2.00
1	1.76
2	1.64
3	1.56
4	1.52
5	1.51
6	1.51



04.4

Complete the sentence.

Choose the answer from the list.

Use TABLE 4. [1 mark]

- 3 minutes
- 4 minutes
- 5 minutes
- 6 minutes

The minimum heating time needed for all of the calcium hydroxide to be changed into calcium oxide and water vapour is



REPEAT OF TABLE 4

Total heating time in minutes	Mass of contents of test tube in grams
0	2.00
1	1.76
2	1.64
3	1.56
4	1.52
5	1.51
6	1.51



Calculate the total mass of water vapour produced by heating the calcium hydroxide.

Use TABLE 4. [2	marks]	
Mass =	q	



The word equation for the reaction is:

calcium oxide + water

The reaction is reversible.

When 4.00 g of calcium hydroxide is completely changed into calcium oxide and water:

- 3.03 g of calcium oxide is produced
- 5.90 kJ of energy is taken in from the surroundings.



0	4	6
	-	

3.03 g of calcium oxide reacts completely with water to produce 4.00 g of calcium hydroxide.

How much energy is transferred to the surroundings in this reaction? [1 mark]

Tick (✓) ONE box.

Less than 5.90 kJ
5.90 kJ
More than 5.90 kJ



0 4.7

The forward reaction takes in energy from the surroundings.

Complete the sentence.

Choose the answer from the list. [1 mark]

- combustion
- endothermic
- exothermic

The forward reaction is

9



0 5
This question is about greenhouse gases and climate change.
05.1
Which TWO gases are greenhouse gases? [2 marks]
Tick (✓) TWO boxes.
Argon
Carbon dioxide
Nitrogen
Methane
Oxygen



0 5.2

Why are greenhouse gases essential for supporting life on Earth? [1 mark]

The percentage of greenhouse gases in the Earth's atmosphere today is increasing.

Many scientists think that this increase is causing global climate change.



0	5	•	3
---	---	---	---

What is a cause of the greenhouse effect?

Complete the sentence. [1 mark]

Greenhouse gases absorb long wavelength



of global

0 5.4
Which TWO are potential effects climate change? [2 marks]

Tick (✓) TWO boxes.

Fewer droughts

Fewer storms

Higher sea levels

Less coastal flooding

Melting polar ice



0	5	5

Water vapour is a greenhouse gas.

The percentage by mass of water vapour in the Earth's atmosphere is 0.25%.

Calculate the mass of water vapour in 350 kg of the Earth's atmosphere.

Give your answer in	J	
Mass =		g



0 6

This question is about fuels.

The energy produced by burning fuels is used to generate electricity in power stations.

TABLE 5, on page 58, shows information about three fuels used to generate electricity.



BLANK PAGE



Į	5)
L		
		_
	Y	
•		
_		_

	FUEL		
	COAL	OIL	NATURAL GAS
State of fuel at room temperature	solid	liquid	gas
Transportation of fuel to power station	train	pipeline	pipeline
Percentage by mass of sulfur in fuel (%)	5		0.001
Relative quantity of solid particles produced when fuel is burned	high	medium	MO



0 6 . 1 Explain

coal is usually transported to power stations by train and NOT by pipeline. Explain why

Use TABLE 5. [2 marks]

le and particulates are atmospheric pollutants produced when fuels are burned. **Sulfur dioxid**

7
9
0

fuel in TABLE 5, on page 58, is burned. 1 kg of each

Which fuel produces the MOST sulfur dioxide?


Give ONE reason for your choice. [2 marks]

Fuel

2	
U	Ŋ
(Q

Give ONE problem caused by sulfur dioxide. [1 mark]

[Turn over]



<u>.</u>

9

<u>..</u> 9

are formed from solid particles. **Particulates**

fuel in TABLE 5, on page 58, is burned. 1 kg of each

Which fuel produces the LEAST particulates?

Give ONE reason for your choice. [2 marks]

Fuel

Reason

6 2

0 6 . 5 Give ON

Give ONE problem caused by particulates. [1 mark]

9 0

Complete the sentence. [1 mark]


Solid particles are formed when fuels undergo incomplete

06.7

FIGURE 6 shows how the use of oil and of natural gas as fuels changed in the UK between 2002 and 2020.

FIGURE 6

Fuel use in the UK in arbitrary units





Describe the trends show [3 marks]	n in FIGURE 6.
[Turn over]	12



0	7

This question is about alloys.

Steels are alloys of iron.

Which non-metal element is in all steels? [1 mark]

Tick (✓) ONE box.

lodine





0	7		2
	-	_	_

Which TWO elements other than iron are in stainless steels? [2 marks]

Tick (✓) TWO boxes.

Chromium

Gold

Magnesium

Nickel

[Turn over]

Zinc



0	7		3
	-	_	

Give TWO properties of stainless steels.

Choose answers from the list. [2 marks]

- brittle
- hard
- low density
- resistant to corrosion
- soluble in water

Property 1	
Property 2	



BLANK PAGE



Titanium is used in alloys.

TABLE 6 shows information about some alloys of titanium.

TABLE 6

TITANIUM ALLOY	M OTHER METALS STRENGTH IN ALLOY		USED IN
	6.0% aluminium	<u></u>	aircraft parts, hip
(4.0% vanadium		joint replacements
۵	5.0% aluminium	\\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\	
۵	2.5% tin	Пдп	all Clait parts
Č	3.0% aluminium		tennis rackets,
)	2.5% vanadium		heart pacemakers



4	•
	<i>σ</i>
	=
2	ס
7:7	
7	-
	り
)
L	n
\$	
8	
	5
\$	
7	פ
4	5
	5
	III de Se
3	
9	1
4	
1	1)
	<u>U</u>

marks]
Σ
6 .
Ш
=
$\overline{\mathbf{m}}$
4
Jse

Mass =

REPEAT OF TABLE 6

TITANIUM ALLOY	MOTHER METALS STRENGTH IN ALLOY		USED IN	
	6.0% aluminium	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	aircraft parts, hip	
(4.0% vanadium		joint replacements	_
	5.0% aluminium	4		
۵	2.5% tin	IIGII	all crait parts	
Č	3.0% aluminium		tennis rackets,	
)	2.5% vanadium		heart pacemakers	






alloy A and alloy B are used to make aircraft parts Suggest why

Use TABLE 6. [1 mark]

10



Titanium alloys used for medical purposes must NOT be toxic. alloy B is NOT used for medical purposes. Suggest why

5, on page 72. [1 mark] **Use TABLE**

BLANK PAGE [Turn over]


0 8

A student investigated the rate of the reaction between zinc and sulfuric acid.

Hydrogen gas is produced during this reaction.

FIGURE 7 shows the apparatus.

FIGURE 7





This is the method used.

- 1. Add 50 cm³ of sulfuric acid to a conical flask.
- 2. Add 2.0 g of zinc to the conical flask.
- 3. Quickly put a stopper in the conical flask and start a timer.
- 4. Measure the time taken to collect 20 cm³ of gas.
- 5. Repeat steps 1 to 4 three more times.

0 8 . 1

Suggest why the stopper must be put in the conical flask as quickly as possible in STEP 3. [1 mark]



The student calculated the rate of the reaction for each trial.

TABLE 7 shows the results of the calculations.

TABLE 7

	TRIAL	TRIAL	TRIAL	TRIAL
	1	2	3	4
Rate of reaction in cm ³ /s	0.78	0.81	0.68	0.81

Determine the mean time taken to collect 20 cm³ of gas.

Do NOT include any anomalous results.



Use the equation:

mean rate of reaction = volume of gas collected mean time taken

[5 marks]			



Mean time taken =	S

08.3

The student changed the investigation so that the mean time taken to collect 20 cm³ of gas was greater.

Which TWO changes would increase the mean time taken to collect 20 cm³ of gas? [2 marks]



Tick (✓) TWO boxes.		
	Use a catalyst	
	Use a larger conical flask	
	Use a lower temperature	
	Use smaller pieces of zinc	
	Use sulfuric acid of a lower concentration	



08.4
Hydrogen gas is produced during this reaction.
Describe the test for hydrogen gas.
Give the result of the test. [2 marks] Test
Result



0 9

This question is about alcohols and carboxylic acids.

Alcohols are used as fuels.

A student burned 1.00 g of six alcohols and determined the energy released from each.

TABLE 8, on page 84, shows the results.



TABLE 8

Alcohol	Formula of one molecule of the alcohol	Energy released in kJ/g
Ethanol	C ₂ H ₅ OH	29.6
Propanol	C ₃ H ₇ OH	33.6
Butanol	C ₄ H ₉ OH	36.1
Pentanol	C ₅ H ₁₁ OH	37.7
Hexanol	C ₆ H ₁₃ OH	38.9
Heptanol	C ₇ H ₁₅ OH	39.8



0	9		1
		_	-

Calculate the mass of ethanol that must be burned to release the same amount of energy as burning 1.00 g of heptanol.

[2 marks]

Mass =	a



REPEAT OF TABLE 8

Alcohol	Formula of one molecule of the alcohol	Energy released in kJ/g
Ethanol	C ₂ H ₅ OH	29.6
Propanol	C ₃ H ₇ OH	33.6
Butanol	C ₄ H ₉ OH	36.1
Pentanol	C ₅ H ₁₁ OH	37.7
Hexanol	C ₆ H ₁₃ OH	38.9
Heptanol	C ₇ H ₁₅ OH	39.8

0 9 . 2


The energy released in kJ/g varies with the number of carbon atoms in one molecule of each alcohol.

Plot the data from TABLE 8 on FIGURE 8, on the opposite page. [2 marks]



FIGURE 8

Energy released in kJ/g



BLANK PAGE



0 9 . 3

Estimate the energy released in kJ when 1.00 g of octanol (C₈H₁₇OH) is burned.

Use FIGURE 8, on page 87. [1 mark]

Energy released = kJ



Carbon dioxide is produced when alcohols are burned.

Carbon dioxide is identified by bubbling the gas through limewater.

Complete the sentence.

Choose the answer from the list. [1 mark]

- calcium chloride
- calcium hydroxide
- calcium nitrate
- calcium sulfate

Limewater is an aqueous solution of



0	9		5
		_	

Give the result of the test when carbon dioxide is bubbled through limewater.

[1 mark]



Ethanoic acid can be produced from ethanol.



What is reacted with ethanol to produce ethanoic acid? [1 mark]

Tick (✓) ONE box.

A halogen
An alkali metal
An oxidising agent
Water



0 9 . 7

Ethanoic acid contains the functional group –COOH

Complete the displayed structural formula of this functional group. [1 mark]

— С О

O - H



0 9 . 8

Ethanoic acid reacts with different compounds.

On the opposite page, draw ONE line from each compound to a product of the reaction of the compound with ethanoic acid. [2 marks]



Compound

Product of the reaction with ethanoic acid

Carbon dioxide

Ethanol

Ethene

Sodium carbonate

Ethyl ethanoate

Hydrogen

Poly(ethene)

[Turn over]



11

1 0

This question is about chemical analysis.

Potassium bromide is used in medicine.

A scientist tested a sample of medicine to show the presence of potassium ions and of bromide ions.

The sample is soluble in water.

10.1

Plan a method the scientist could use to show that the sample of medicine contains potassium ions AND bromide ions.



			4 -	4	
I	CCI	n	ITIC	•	has:
	36			L	Hab.

- a Bunsen burner
- a metal wire
- test tubes
- a dropping pipette
- distilled water
- dilute nitric acid
- silver nitrate solution.

[6 marks]					







100

The scientist could also use an instrumental method to show the presence of potassium ions in the medicine.

10.2

Which instrumental method could be used to show the presence of potassium ions in the medicine? [1 mark]



1	0		3
_	_	_	

Give ONE advantage of using this instrumental method instead of a chemical test. [1 mark]

END OF QUESTIONS

8



Additional page, if required.		
Write the question numbers in the left-hand margin.		



Additional page, if required.		
Write the question numbers in the left-hand margin.		



BLANK PAGE

For Examiner's Use			
Question	Mark		
1			
2			
3			
4			
5			
6			
7			
8			
9			
10			
TOTAL			

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 AQA and its licensors. All rights reserved.

WP/M/SB/Jun23/8462/2F/E3



