



# **Teacher Support Materials**

## **Maths GCE**

### **Paper Reference MFP1**

Copyright © 2008 AQA and its licensors. All rights reserved.

Permission to reproduce all copyrighted material has been applied for. In some cases, efforts to contact copyright holders have been unsuccessful and AQA will be happy to rectify any omissions if notified.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334). Registered address: AQA, Devas Street, Manchester M15 6EX.  
*Dr Michael Cresswell*, Director General.

## Question 1

1 The matrices  $\mathbf{A}$  and  $\mathbf{B}$  are given by

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 3 & 8 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

The matrix  $\mathbf{M} = \mathbf{A} - 2\mathbf{B}$ .

(a) Show that  $\mathbf{M} = n \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$ , where  $n$  is a positive integer. (2 marks)

(b) The matrix  $\mathbf{M}$  represents a combination of an enlargement of scale factor  $p$  and a reflection in a line  $L$ . State the value of  $p$  and write down the equation of  $L$ . (2 marks)

(c) Show that

$$\mathbf{M}^2 = q\mathbf{I}$$

where  $q$  is an integer and  $\mathbf{I}$  is the  $2 \times 2$  identity matrix. (2 marks)

## Student Response

1a)	$\mathbf{A} - 2\mathbf{B} = \begin{bmatrix} 2 & 1 \\ 3 & 8 \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 6 & 8 \end{bmatrix} = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix}$
	$\begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix} = 3 \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$ ✓
b)	$p=3$ ✓ $L =$ reflection along line $y=x$ . ✗
c)	$\mathbf{M}^2 = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix} \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix} = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix}$
	$\begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix} = 9 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ✓

## Commentary

This question was generally very well answered. The most common mistake was in part (b), as exemplified in the chosen script, where the mirror line was given as  $y = x$  rather than  $y = -x$ . It was, however, acceptable to give this reflection in conjunction with an enlargement with scale factor  $-3$ .

**Mark scheme**

Q	Solution	Mark	Total	Comments
1(a)	$M = \begin{bmatrix} 0 & -3 \\ -3 & 0 \end{bmatrix}$	B2,1	2	B1 if subtracted the wrong way round
(b)	$p = 3$	B1F		ft after B1 in (a)
	$L$ is $y = -x$	B1	2	Allow $p = -3$ , $L$ is $y = x$
(c)	$M^2 = \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix}$	B1F		Or by geometrical reasoning; ft as before
	$\dots = 9I$	B1F	2	ft as before
	<b>Total</b>		<b>6</b>	

## Question 2

- 2 (a) Show that the equation

$$x^3 + x - 7 = 0$$

has a root between 1.6 and 1.8.

(3 marks)

- (b) Use interval bisection twice, starting with the interval in part (a), to give this root to one decimal place.

(4 marks)

## Student response

2(a)	Let $f(x) = x^3 + x - 7$ <del><math>f(1.6) = 1.6^3 + 1.6 - 7 = -1.304 &lt; 0</math></del> $f(1.6) = 1.6^3 + 1.6 - 7 = -1.304 < 0$ $f(1.8) = 1.8^3 + 1.8 - 7 = 0.632 > 0$ ✓ change of sign ∴ the equation $x^3 + x - 7$ has a root between 1.6 and 1.8 ✓	Leave blank
(b)	$\frac{1.6 + 1.8}{2} = 1.7$ $f(1.7) = 1.7^3 + 1.7 - 7 = -0.387$ $-0.387 < 0$ $\frac{1.7 + 1.8}{2} = 1.75$ $f(1.75) = 1.75^3 + 1.75 - 7$ $= 0.109375$ $\approx 0.1$ (1 d.p.) ✓	3 3 0 <hr/> 6

## Commentary

Almost all the candidates were able to make a good start to this question, though some failed to draw a proper conclusion in part (a). In part (b) the response was again very good. The most common error was a failure to grasp what exactly was being asked for at the end of the question. If it is known that the root lies between 1.7 and 1.75, then it must be 1.7 to one place of decimals. But many candidates gave no value to 1 DP, or, like this candidate, gave a value of the function rather than a value of  $x$ .

## Mark Scheme

2(a)	$f(1.6) = -1.304$ , $f(1.8) = 0.632$ Sign change, so root between	B1,B1 E1	3	Allow 1 dp throughout
(b)	$f(1.7)$ considered first $f(1.7) = -0.387$ , so root $> 1.7$ $f(1.75) = 0.109375$ , so root $\approx 1.7$	M1 A1 m1A1	4	m1 for $f(1.65)$ after error
	<b>Total</b>		7	

## Question 3

3 It is given that  $z = x + iy$ , where  $x$  and  $y$  are real numbers.

(a) Find, in terms of  $x$  and  $y$ , the real and imaginary parts of

$$z - 3iz^*$$

where  $z^*$  is the complex conjugate of  $z$ .

(3 marks)

(b) Find the complex number  $z$  such that

$$z - 3iz^* = 16$$

(3 marks)

## Student Response

3a) $z - 3iz^*$	
$x + iy - 3i(x - iy)$	
$x + iy - 3ix - 3i^2y$ X	
$x + iy - 3ix + 3y$	
Real = $x + 3y$ ✓	2
Imaginary = $y - 3x$	
b) $z - 3iz^* = 16$	
$x + iy - 3ix + 3y = 16$	
$x + 3y = 16$	
$-9x + 3y = 0$	
$10x = 16$ ✓	$z = 1.6 + 4.8i$ ✓
$x = 1.6$ ✓	
$y = 4.8$ ✓	3
	<u>5</u>

## Commentary

Although most of the candidates who take this paper are good at algebra, there are still quite a number who make elementary sign errors. This candidate is typical of many who made the same error, affecting the real part of the given complex number. The candidate then uses the incorrect expression in part (b), but does correct work to arrive at a value of  $z$ . Full marks were awarded in part (b) for candidates who had lost a mark in part (a) for this particular error.

## Mark Scheme

<p><b>3(a)</b></p>	<p>Use of <math>z^* = x - iy</math>  <math>z - 3iz^* = x + iy - 3ix - 3y</math>  <math>R = x - 3y, I = -3x + y</math></p>	<p>M1  m1  A1</p>	<p>3</p>	<p>Condone sign error here  Condone inclusion of <math>i</math> in <math>I</math>  Allow if correct in (b)</p>
<p><b>(b)</b></p>	<p><math>x - 3y = 16, -3x + y = 0</math>  Elimination of <math>x</math> or <math>y</math>  <math>z = -2 - 6i</math></p>	<p>M1  m1  A1F</p>	<p>3</p>	<p>Accept <math>x = -2, y = -6</math>;  ft <math>x + 3y</math> for <math>x - 3y</math></p>
<b>Total</b>			<p><b>6</b></p>	